

## State of Science

# Geomorphology under ice streams: Moving from form to process

Chris R. Stokes\* 

Department of Geography, Durham University, Durham DH1 3LE, UK

Received 27 March 2017; Revised 1 September 2017; Accepted 7 September 2017

\*Correspondence to: Chris R. Stokes, Department of Geography, Durham University, Durham, DH1 3LE, UK. E-mail: c.r.stokes@durham.ac.uk

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

# ESPL

Earth Surface Processes and Landforms

**ABSTRACT:** Ice streams are integral components of an ice sheet's mass balance and directly impact on sea level. Their flow is governed by processes at the ice-bed interface which create landforms that, in turn, modulate ice stream dynamics through their influence on bed topography and basal shear stresses. Thus, ice stream geomorphology is critical to understanding and modelling ice streams and ice sheet dynamics. This paper reviews developments in our understanding of ice stream geomorphology from a historical perspective, with a focus on the extent to which studies of modern and palaeo-ice streams have converged to take us from a position of near-complete ignorance to a detailed understanding of their bed morphology. During the 1970s and 1980s, our knowledge was limited and largely gleaned from geophysical investigations of modern ice stream beds in Antarctica. Very few palaeo-ice streams had been identified with any confidence. During the 1990s, however, glacial geomorphologists began to recognise their distinctive geomorphology, which included distinct patterns of highly elongated mega-scale glacial lineations, ice stream shear margin moraines, and major sedimentary depocentres. However, studying relict features could say little about the time-scales over which this geomorphology evolved and under what glaciological conditions. This began to be addressed in the early 2000s, through continued efforts to scrutinise modern ice stream beds at higher resolution, but our current understanding of how landforms relate to processes remains subject to large uncertainties, particularly in relation to the mechanisms and time-scales of sediment erosion, transport and deposition, and how these lead to the growth and decay of subglacial bedforms. This represents the next key challenge and will require even closer cooperation between glaciology, glacial geomorphology, sedimentology, and numerical modelling, together with more sophisticated methods to quantify and analyse the anticipated growth of geomorphological data from beneath active ice streams. © 2017 The Authors. Earth Surface Processes and Landforms published by John Wiley & Sons Ltd.

**KEYWORDS:** ice stream; glacial geomorphology; West Antarctica; subglacial bedforms; subglacial sediments

## Introduction

An ice stream is defined as 'a region in a grounded ice sheet in which the ice flows much faster than in the regions on either side' (Paterson, 1994: 301). They are large features (1–10 s of km wide and 10–100 s of km long) and their significance arises from their high velocity (100 s–1000 s  $\text{m a}^{-1}$ ) and their correspondingly large ice flux. In Greenland, for example, ice streams account for approximately 50% of the ice sheet's annual mass loss (van den Broeke *et al.*, 2009), and this figure increases to around 90% in Antarctica (Bamber *et al.*, 2000), where there is much less surface melt (the other major process through which ice sheets lose mass). Thus, the dynamics of ice streams holds important implications for ice sheet mass balance and associated impacts on global sea level, which has become an increasingly serious concern since the 1970s (Hughes, 1973; Mercer, 1978; Alley *et al.*, 2005; Shepherd *et al.*, 2012; Hanna *et al.*, 2013; Joughin *et al.*, 2014). Although there are uncertainties about the precise atmospheric and oceanic

factors that force increased discharge via ice streams (Carr *et al.*, 2013) and the extent to which subglacial topography modulates their response (Pfeffer *et al.*, 2008; Nick *et al.*, 2013; Favier *et al.*, 2014; Morlighem *et al.*, 2014), their rapid flow is largely governed by processes at the ice-bed interface, which act to sustain, enhance or inhibit their motion (see reviews in Bentley, 1987; Bennett, 2003; Stokes *et al.*, 2007). Explicit comparisons have rarely been made, but the ice-bed interface is analogous to the 'boundary layer' in fluvial and aeolian environments, where shear stresses (both basal and lateral in the case of ice streams) oppose the flow of the overlying medium. This analogy extends further because processes within the boundary layer create a distinctive geomorphology that is characterised by subglacial bedforms that resemble features created in fluvial and aeolian environments (Shaw 1983; Fisher and Shaw, 1992; Shaw *et al.*, 2008; Clark, 2010; Barchyn *et al.*, 2016; Ely *et al.*, 2016).

The study of ice stream geomorphology is, therefore, critical to our understanding of ice stream behaviour. The creation of

landforms beneath ice streams results from erosion, transport and deposition of sediment which, in turn, is intimately linked to the mechanisms through which they are able to flow so rapidly (Ó Cofaigh *et al.*, 2002; Schoof, 2002; Alley *et al.*, 2003; Bingham and Siegert, 2009; Graham *et al.*, 2009). Over short time-scales (years to decades), the evolution of subglacial geomorphology influences the roughness of the bed and the routing and pressure of subglacial meltwater, which further impacts on bed 'stickiness' and modulates the flow of the ice stream (Schoof, 2002). Over longer time-scales (centuries to millennia), these processes may alter the subglacial morphology in a more gradual manner, such as over-deepening the bed (Alley *et al.*, 2003; Cook and Swift, 2012) and creating major sediment depocentres at their terminus (Alley *et al.*, 2007). Unlike fluvial and aeolian environments, however, the basal environment of ice streams is notoriously difficult to access and observe, and there is far more uncertainty in securely linking form to process. This, in turn, severely limits attempts to understand ice stream behaviour and parameterise their basal conditions in numerical modelling (Bentley, 1987; Bennett, 2003; Ritz *et al.*, 2015).

Despite the difficulty in observing the subglacial environment, our understanding of ice stream geomorphology has grown rapidly in the last three decades, from almost complete ignorance to a detailed knowledge of their geomorphological imprint. This has been brought about by two main approaches: (i) borehole and geophysical investigation of modern (active) ice streams; and (ii) sedimentological, geomorphological and geophysical investigation of well-preserved palaeo-ice stream beds. The aim of this paper is to comprehensively review progress in these two main areas from a historical perspective; highlight the key questions that remain; and discuss the opportunities that are likely to arise that will enable them to be addressed. The following section reviews the progress made through the pioneering geophysical and borehole investigations on active ice streams, mostly in West Antarctica, from the 1970s to the early 2000s. The third section reviews the parallel progress made by those seeking to identify and characterise the beds of palaeo-ice sheets from the late 1980s to early 2000s. The fourth section returns to a more recent body of work aimed at imaging the beds of active ice streams (generally post-2000), and highlights the links between these geophysical investigations of modern ice streams and the large growth in data from palaeo-ice stream beds. This convergence of knowledge has led to some important advances in our understanding of the geomorphology of ice stream beds, but it remains a key challenge to securely link landforms to processes. The fifth section focuses on highlighting the key questions that remain and briefly highlights opportunities that might arise to address them. In the final section it is concluded that future surveys of modern-ice stream beds are likely to offer unprecedented opportunities to study ice stream geomorphology that will require the continued cross-pollination of ideas and concepts from each of the two main approaches, together with a third approach – numerical modelling.

## Early Exploration of the Beds of Active Ice Streams

### The discovery and definition of ice streams


The first use of the term 'ice stream' can be traced back to Rink (1877), who used 'Is-Ström' (i.e. 'ice stream' or 'ice current') in relation to the Greenland Ice Sheet (GrIS) to describe 'those outer parts of the inland ice which are moving with greater

rapidity towards the ice-fjords than the rest of its outer margin' (p. 369). Almost two decades on, Bell (1895) used 'ice stream' to describe flow in the vicinity of Hudson Strait, which is the first use of the term in relation to a palaeo-ice sheet (Brookes, 2007). In a paper entitled 'Antarctica and Some of its Problems', Edgeworth (1914) appears to be the first to use the term in relation to that region, noting that ice stream 'differ from ordinary glaciers in that they are not bounded by rock walls, but are simply slight depressions in the general surface of the inland ice, marking areas where the ice is in more rapid movement than in adjacent areas' (p. 611). Thus, whilst the definition of an ice stream is often attributed (Bentley, 1987; Bennett, 2003) to a note by Charles Swithinbank (Swithinbank, 1954), he was simply stating a preference for the term used by Rink (1877) and Edgeworth (1914) over other possibilities 'on account of the impression it conveys of movement' and because 'it was the term first used in describing these features' (p. 186; see also the note by Roscoe (1954) in the same issue).

The early descriptions and discussions of ice streams (Rink, 1877; Edgeworth, 1914; Swithinbank, 1954) emphasised that they have no exposed rock to define their lateral margins, thereby distinguishing them from outlet glaciers. However, adhering to such a strict definition is often deemed impractical (cf. Bentley, 1987; Paterson, 1994), not least because the vast majority of ice streams *sensu stricto* become bordered by rock-walls in coastal regions, where they flow through deep fjords, or may lie within deep glacial valleys where rock walls lie just beneath the ice surface. This is not the case for a small number of present-day ice streams that flow over relatively flat terrain along the Siple Coast of West Antarctica, and for several ice streams that operated at the land-terminating margins of palaeo-ice sheets (Margold, Stokes, Clark, and Kleman, 2015b). Thus, a more useful distinction from a process point of view is to define ice streams according to their topographic setting (Bentley 1987). Following Bentley (1987), Stokes and Clark (1999) used the term 'pure ice streams' (see also Paterson, 1994) for those not associated with large troughs, and 'topographic ice streams' for those that are constrained by topography. Truffer and Echelmeyer (2003) used the terms 'ice stream' (for 'pure') and 'isbræ' (for topographic) to describe these two types, and pointed out important differences in their geometry, flow mechanisms, force balance, hydrology and potential for instability. However, they also emphasised (as did Bentley, 1987) that these two types are end members along a spectrum, with many ice streams displaying characteristics of both types (see Table 1 in Truffer and Echelmeyer, 2003). For example, enhanced flow within the tributaries of an ice stream might be strongly steered by topography, even if the lower reaches of the main ice stream 'trunk' is not, as has been suggested for ice streams along the Siple Coast (Joughin *et al.*, 1999). In this paper, I will refer to ice streams in the broadest sense and include fast flowing outlet glaciers that are topographically controlled (cf. McIntyre, 1985; Bentley, 1987; Paterson, 1994).

### West Antarctic ice streams and 'glaciology's grand unsolved problem'

Given their early recognition (e.g. Rink, 1877) and now well-established significance (e.g. Bennett, 2003), there were very few studies of ice streams until the 1970s, see Figure 1. Indeed, following Swithinbank's (1954) brief note, the first scientific paper with 'ice stream' or 'ice streams' in its title was Hughes' (1977) review of 'West Antarctic Ice Streams', but it is important to note that several major outlet glaciers (ice streams in the broadest sense) had been studied, e.g. Lambert Glacier in East



**Figure 1.** The number of papers with 'ice stream' or 'ice streams' in the title from the ISI Web of Science catalogue ( $n = 538$ ; search date: 14 March 2017). The paper from 1976 is a conference abstract from the Midwestern AGU meeting (Bentley, 1976) and so the first major paper was by Hughes (1977). A slow growth in outputs covering ice streams only begins in the mid-1980s, following the pioneering geophysical studies aimed at characterising their subglacial environment (Alley *et al.*, 1986; Blankenship *et al.*, 1986). Indeed, the anomalous peak in 1987 is due to a special issue (SI) of the *Journal of Geophysical Research* that contained papers from the Chapman Conference on 'Fast Glacier Flow: Ice Streams, Surging and Tide-water Glaciers' (see Clarke, 1987). A further notable peak results from a collection of the 'Siple Coast Project' papers that were published together in *Journal of Glaciology* in 1993 (vol. 133). The most productive year remains 2003, which coincides with the production of two thematic issues related to ice streams: the 'International Symposium on Fast Glacier Flow' in the *Annals of Glaciology* and the 'Palaeo-Ice Stream International Symposium' in *Boreas* in 2003. Note that the focus here is on the emergence of the term 'ice stream' in scientific papers and that some studies of ice streams/outlet glaciers will be missing (those on Pine Island and Thwaites Glacier, W. Antarctica; Lambert Glacier in East Antarctica; and Jakobshavn Isbræ, in W. Greenland), in addition to some edited book chapters.

Antarctica (Morgan and Budd, 1975; Allison, 1979). Ice streams had, of course, been recognised for some time, but mechanisms of fast glacier flow had been investigated almost exclusively in relation to surge-type glaciers (Meier and Post, 1969). Indeed, in a comment published at the end of Meier and Post's (1969) classic paper ('What are glacier surges?'), Johannes Weertman was concerned that they had ignored fast-moving outlet glaciers (i.e. ice streams *sensu lato*), suggesting that their flow mechanisms may actually be similar to surging glaciers, and imploring the community not to over-look the 'outlet glacier problem' (Weertman, 1969: p. 817). With hindsight, it is clear that the observations, concepts and theories that had been applied to surging glaciers would turn out to be instrumental in shaping much of the early work on ice streams, not least because many of those who had studied surging glaciers began to turn their attention to ice streams.

It was concern over the stability of the marine-based West Antarctic Ice Sheet (WAIS) that instigated a concerted effort to understand its dynamics and stability. Hughes (1972, 1973) was one of the first to hypothesise that the junction of grounded ice sheet with a floating ice shelf (the grounding line) might be unstable if the bed deepens inland. He pointed out that this was the case for the WAIS (Hughes, 1972, 1973) and went so far as to suggest that retreat was already underway. A further analytical treatment by Weertman (1974) confirmed the basic idea of marine-ice sheet instability, but he was more equivocal in stating whether or not the WAIS was already undergoing marine ice sheet instability (Joughin *et al.*, 2014 and Favier *et al.*, 2014 for more recent analyses of this important issue). Weertman (1974) acknowledged the possibility that WAIS was disintegrating, but also noted that a complete treatment of this problem must take into account the fast moving ice streams that exist along the Siple Coast, where it flows into the Ross Ice Shelf. Later work by Hughes (1977) argued that the ice streams

themselves were inherently unstable, but he acknowledged that the factors driving their motion were not well understood and highlighted the need for further observations, which he suggested should include bed profiling using radar and seismic surveys, and borehole investigations. The need for these investigations was made more urgent by a series of high-profile papers that debated the potential instability of the WAIS (Thomas, 1976; Weertman, 1976; Whillans, 1976; Mercer, 1978; Thomas *et al.*, 1979), and which Weertman (1976) referred to as 'Glaciology's grand unsolved problem'.

Hughes' (1977) invocation for geophysical exploration of ice streams was, in fact, already underway as a result of collaboration between the UK's Scott Polar Research Institute (SPRI) and the Technical University of Denmark (TUD), with the cooperation and support of the National Science Foundation (NSF). Airborne radio echo-sounding of a 50 km grid in Marie Byrd Land was undertaken, where the Siple Coast ice streams (then named A, B, C, D and E)<sup>1</sup> had earlier been reported based on reconnaissance flights using the same technique (Robin *et al.*, 1970). The results were reported in Rose (1979), which was one of the first studies to investigate the basal conditions beneath active ice streams, together with Morgan and Budd's (1975) radio-echo sounding of the Lambert Glacier basin. A key conclusion from Rose's (1979) study was that the lower reaches of the ice streams were contained within shallow, broad channels within a relatively smooth topography that becomes progressively rougher in their upstream reaches. Rose also calculated basal shear stresses using ice surface slopes and thicknesses and found that they were less than 15 kPa close to the grounding line. His first-order calculations of basal

<sup>1</sup>These ice streams have since been re-named Mercer (A), Whillans (B), Kamb (C), Bindschadler (D) and MacAyeal (E), but the original names are used here to aid cross-referencing to the older literature.

temperatures confirmed that the beds of the ice streams were warm-based, whereas the inter-stream ridges were below the pressure melting point. Similar conclusions were reached by Morgan and Budd (1975), and by Stephenson and Doake (1982), which was one of the first geophysical studies of Rutford Ice Stream, West Antarctica, and which highlighted the sensitivity of its grounding line position to changes in ice thickness. Crabtree and Doake (1982) also used radio-echo sounding to provide some of the first data from the bed of Pine Island Glacier and noted its potential vulnerability to rapid retreat from a bedrock sill, a topic which has attracted considerable attention in recent years (Favier *et al.*, 2014; Smith *et al.*, 2017). Rose (1979) also pointed out the enigmatic behaviour of Ice Stream C, which appeared to be inactive, but displayed many of the surficial and internal characteristics of the neighbouring active ice streams.


Another key conclusion from Rose's (1979) investigation was that the flow mechanism of pure ice streams was fundamentally different from that of normal ice sheet flow. Similar arguments were put forward by McIntyre (1985) who noted the difference in flow mechanisms between major ice streams/outlet glaciers that occupied deep topographic troughs (and sometimes coincide with topographic steps) and those along the Siple Coast, which he suggested might have more in common with ice shelves. Notwithstanding the enigmatic Ice Stream C, therefore, a key question was how these particular ice streams achieved their high velocities, given the extremely low ice surface slopes and low driving stresses. This prompted a series of pioneering geophysical investigations aimed at characterising their basal environment and ascertaining the mechanism(s) of flow.

### Ice stream geophysics: A paradigm shift in glaciology?

In 1983, a major field programme was initiated to study the Siple Coast ice streams and geophysical techniques targeted one of the two major tributaries of Ice Stream B at the 'Upstream B camp' (UpB), where ice thicknesses approach 1000 m and

velocities exceed  $400 \text{ m a}^{-1}$  (Whillans *et al.*, 1987). The field survey during the first season of the programme (1983–1984) deployed seismic reflection techniques to investigate the nature of the contact zone between the ice stream and the bed, and the results were presented in two papers (Alley *et al.*, 1986; Blankenship *et al.*, 1986), that Clarke (1987) soon suggested would resonate 'for many years' (p. 8840). These surveys revealed a layer of sediment averaging 5–6 m thick immediately beneath the ice with very low compressional (P) and shear (S) wave speeds, see Figure 2 (Blankenship *et al.*, 1986). Such low wave speeds were taken to imply that the sediment was highly porous and saturated with water, and was too weak to support the shear stress exerted by the overlying ice. Further arguments relating to till porosity, the ice stream force balance, and the water balance were presented in the accompanying paper by Alley *et al.* (1986), who argued that 'deformation within the till is the primary mechanism by which the ice stream moves' (p. 57).

Boulton's (1986) commentary that appeared alongside these papers hailed them as a 'paradigm shift' in glaciology because most previous models of glacier flow had assumed that ice moved over a rigid surface. However, he also emphasised that a 'rigid bed model' would have seemed quite implausible to glacial geologists who were familiar with the sediments and landforms on the beds of palaeo-ice sheets, noting that its survival for so long 'is an indication of how little note the glaciological community took of the views of geologists' (p. 18). Indeed, Boulton pointed out that the discovery of deformable sediments beneath these ice streams was entirely consistent with what had long been known from investigations on palaeo-ice sheet beds, such as glacially induced deformation structures (Slater, 1926) that are often associated with subglacial bedforms, such as drumlins (Stanford and Mickelson, 1985). Moreover, Boulton and Jones (1979) had earlier presented experimental evidence of till deforming beneath Breiðamerkurjökull in Iceland, the results of which received a more thorough treatment in the classic paper by Boulton and Hindmarsh (1987), which also described the geological consequences in terms of the development of



**Figure 2.** One of the first glimpses of the geomorphology of an active ice stream bed, from the UpB site on Ice Stream B, West Antarctica (from Blankenship *et al.*, 1986). Blankenship *et al.* (1986) noted that flat, parallel reflectors in the along-flow direction (a) contrasted with those observed in the across-flow direction (b). Note the 'hump' in one of the reflector just beneath the ice bottom (i.e. buried within the till layer) around 800 m along the profile, labelled 'layer bottom (?)' on panel (b). Reproduced by permission of Nature Publishing Group. [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]

subglacial bedforms. Boulton's (1986) commentary ended with a plea for glaciologists and glacial geologists to work together on coupling the form and structure of ice sheet beds with ice dynamics 'to the benefit of both disciplines, in a way which until now has been sadly lacking' (p. 18).

It is often overlooked that both Blankenship *et al.* (1986) and Alley *et al.* (1986) speculated about buried features within the till layer, albeit from observations with limited resolution (Figure 2), and complicated by acquisition artefacts such as hyperbolae (Rooney, 1987a). Blankenship *et al.* (1986) noted how the occurrence of more than one reflector suggested some stratification within the till layer, which was nearly flat in the along-flow direction. In contrast, there was much less continuity in the reflectors in the across-flow direction, and they noted a 'hump' that seemed to rise near, or even to, the base of the ice (Figure 2), with another layer-bottom reflector indicating localised relief of 8 m. They described the emerging picture as 'one of ridges and valleys in the basal surface of the subglacial layer, trending parallel to the axis of the ice stream' with their alignment suggesting 'they are formed by erosion' (p. 56). Alley *et al.* (1986) went further and described the features as flutes (citing Gravenor and Meneley, 1958), suggesting they may be a characteristic erosional consequence of deforming till, and outlining further implications of their erosional-deformational model. They argued that variations in till thickness over the flutes should cause fluctuations in basal drag that affect ice stream flow, with higher drag over the ridge crests. A further corollary is that the estimated till flux at UpB is equivalent to steady state erosion of  $\sim 0.5 \text{ mm a}^{-1}$  over the catchment, which would eventually be transported and delivered to the grounding line to form a 'till delta' (Alley *et al.*, 1986).

An AGU Chapman Conference on Fast Glacier Flow in 1986 provided an opportunity for several workers to discuss the pioneering discoveries from under Ice Stream B (see introduction by Clarke, 1987), and a collection of papers were published in a special issue of the *Journal of Geophysical Research* in 1987 (Figure 1). However, only a handful of these papers discussed the geomorphology of the ice-bed interface, and none in any detail. Indeed, Bentley's (1987) influential review in that special issue simply reiterated Hughes' (1977) earlier plea for more data on the bed conditions, emphasising that all of the existing numerical models of ice stream flow (except for Alley, 1989b, also in the same issue) suffer from a lack of measurements of the physical conditions at the bed and the lack of a valid sliding law. Shabtaie *et al.* (1987) described the morphology of Ice Streams A, B and C from airborne radar sounding and over-snow traverses, but their focus was primarily on the ice surface morphology. However, they were able to confirm many of the underlying topographic features reported by Rose (1979) and they also detected localised patches of strong reflectors at the bed of Ice Stream C, which they took to imply the existence of a water layer at least several centimetres to several metres thick, and which was probably ponded in some locations.

A series of four papers expanded on the earlier geophysical investigations at UpB Camp. Blankenship *et al.* (1987) presented more detailed results of one of the active seismic measurements, calculating that the porosity of the till was greater than 0.32 and probably closer to 0.4, and was saturated with water at a pore pressure only about 50 kPa less than the ice overburden pressure (9000 kPa). The second paper by Rooney *et al.* (1987b) offered greater insight into the characteristics of the buried features within the till layer in this location. They reported data from two parallel seismic reflection lines that were 8.3 km long and oriented perpendicular to ice flow and 360 m apart. These lines revealed a near-continuous till layer averaging  $6.5 \pm 0.5 \text{ m}$  and characterised by an undulating surface of ridges and troughs aligned parallel to ice flow at the

bottom of the till layer. They considered whether the largest of these features may have been a drumlin, but suggested that the dimensions were not typical of drumlins. Rather, and following Alley *et al.* (1986), they argued that these buried features (200–1000 m wide and 6–13 m deep) were similar to flutes observed on the bed of the Laurentide Ice Sheet (LIS) (citing Gravenor and Meneley, 1958, and Smith, 1948). Interestingly, Smith (1948) preferred the term 'giant glacial grooves' for the features he observed along the Mackenzie Valley in north-western Canada, and which extend up to 13 km in length (see section 'Giant glacial grooves and mega-scale glacial lineations').

Rooney *et al.* (1987a) also drew attention to the base of the till layer, where a gently dipping sedimentary structure (interpreted to be sedimentary bedrock: see also Rooney *et al.*, 1987b) was truncated by the till in an angular unconformity that suggested that the till was eroding, or had eroded, its bed. Indeed, they pointed to disturbed zones of anomalous dip direction immediately beneath the till that might represent the fracture and entrainment of large bedrock rafts into the till layer or deeper sub-sole deformation structures some 30 m below the ice-bed interface. In the third paper, Alley (1989a) presented further evidence to argue that the till beneath Ice Stream B was deforming, eroding its bed, and being deposited at the grounding line as till deltas tens of kilometres long (see also Alley *et al.*, 1989a). They pointed out that evidence for such deltas may already have been found in the grounding line mapping of Shabtaie and Bentley (1987), which showed lobate regions of grounded ice tens of kilometres long at the mouths of Ice Stream's A and B. Such deltas might be particularly important for ice dynamics because their deposition serves to stabilise the grounding line by locally increasing the water depth required for flotation (see later work by Alley *et al.*, 2007). A further implication was that till deltas should mark recessional positions of marine-based ice sheets. The fourth paper (Alley, 1989b) developed a simple one-dimensional numerical model of ice stream flow on deforming till with a linear viscous rheology, although the possibility of partial decoupling across a water film was considered and, they argued, was likely to be proportionally more significant closer to the grounding line.

An important theme from the collection of papers on 'Fast Glacier Flow' (Clarke, 1987) was that the geometry and pressure of the basal water system was 'very important in facilitating rapid basal velocities' (Alley, 1989a: p. 110). Irrespective of whether the ice slid over a till layer or deformed it, rapid motion would only occur if the water pressure was relatively close to the ice overburden pressure (Boulton and Hindmarsh, 1987; Clarke, 1987). In turn, the water pressure depends on the water supply and its drainage path, which is likely to be influenced by the bed geomorphology and may also play a role in shaping the form of the bed (e.g. through erosion and deposition). In relation to Ice Stream B, it had been argued that excess subglacial water was incapable of being conducted through subglacial aquifers (Lingle and Brown, 1987) or advected within a deforming bed (Alley, 1989a), and so it was assumed that it drained at the ice-bed interface. However, the form of that basal drainage system was largely unknown. Alley and co-workers developed a series of hypotheses regarding the water-drainage system that might be associated with a deforming till layer and which could be tested by direct borehole drilling to the ice-bed interface (Alley, 1989a, b). The major conclusions were that most of the ice stream motion should arise from bed deformation; that most of the subglacial water flowed in a distributed system approximating a thin film at the ice-bed interface (with local thickenings up to a few mm, but with no channels or linked cavities); and that effective

pressures were small and should decrease down the ice stream (Alley 1989a). In terms of the geomorphology, and in the absence of channelized meltwater supplies (e.g. from supraglacial sources), they argued that it was unlikely that ice streams underlain by a metres-thick layer of saturated sediment would host any channelized drainage systems, such as those which might produce discrete channels eroded into the till, because sediment within the till layer would creep rapidly into incipient channels or cavities with lower pressures (Boulton and Hindmarsh, 1987; Alley, 1989a). It was generally assumed, therefore, that the subglacial drainage system would remain distributed and would erode and transport relatively small quantities ( $<0.1 \text{ m}^3 \text{ m}^{-1} \text{ a}^{-1}$ ) of fine-grained sediment (Alley *et al.*, 1989a), and effectively doing very little geomorphological 'work'. Elsewhere, Doake *et al.* (1987) imaged the bed of Rutford Ice Stream in the vicinity of the grounding line and showed that the ice stream was intermittently grounded on raised areas of the bed that formed 'esker-like' features (Doake *et al.*, 1987: p. 8951).

### Direct borehole access: Sliding versus deformation or ploughing

Despite the paradigm shift in understanding brought about by geophysical investigations, Alley *et al.* (1989b) acknowledged that the details of glacier motion on a deforming bed remained elusive and suggested that 'bore-hole observations undoubtedly will be necessary to solve the problems of till deformation beneath thick ice' (p. 137). Indeed, the series of papers published by Alley in 1989 (Alley, 1989a, b; Alley *et al.*, 1989b) were partly motivated by the knowledge that Barclay Kamb and Herman Engelhardt (and co-workers) would soon be drilling to the base of Ice Stream B (Alley, 1989a). The initial results were reported in Engelhardt *et al.* (1990) who were the first to describe the physical conditions at the base of a fast moving Antarctic ice stream. Similar work was also being undertaken at Jakobshavns Isbræ in West Greenland, with the results published in Iken *et al.* (1993), also discussed below.

Using a hot water method, Engelhardt *et al.* (1990) drilled several boreholes to the base of Ice Stream B and confirmed that it was at the pressure melting point (as first calculated by Rose, 1979) and that basal water pressure was within about 160 kPa bars of the ice overburden pressure (broadly similar to the values calculated by Blankenship *et al.*, 1987). Moreover, when the boreholes reached the bed, the water levels within them dropped very rapidly, revealing a connection to a basal water conduit system. Engelhardt *et al.* (1990) sampled some of the basal sediments driven into suspension and, through the use of a borehole penetrometer and piston corer, were able to sample a muddy, unsorted mixture with rock clasts as large as 5 cm, thus resembling a basal till. Weight loss on drying confirmed the relatively high porosity (~40%) calculated from earlier indirect methods (Blankenship *et al.*, 1987). The high porosity, together with a mixed age of recovered microfossils and bedrock sources would be expected from a till that was deforming, but the discovery of a well-connected basal water system at high pressure led Engelhardt *et al.* (1990) to conclude that 'the proportions of ice stream motion due to basal sliding, subglacial till deformation and ice superplasticity remain unknown' (p. 59).

Although it did not contradict the earlier work based on geophysics, the borehole data from Ice Stream B and other non-ice stream settings (Blake *et al.*, 1992; Humphrey *et al.*, 1993; Iverson *et al.*, 1994, 1995) had begun to trigger a re-evaluation of the deforming bed mechanism and, in particular, question the linear viscous rheology of the saturated till

(Murray, 1997). Crucially, it allowed workers to conduct experiments on the till recovered from beneath Ice Stream B and, building on concepts and experiments on the mechanics of clay-rich soils, Kamb (1991) tested freshly-cored till from Ice Stream B using a shear box method. Kamb found that the till sheared rapidly and underwent 'catastrophic' failure that resembled a plastic material with a yield stress of around 2 kPa. This suggested that the flow law for the till was highly non-linear and Kamb presented further arguments that the deforming bed mechanism would be unstable if a linear viscous rheology was assumed, due to the feedback from the generation of basal water by shear heating in basal till (i.e. the ice stream should rapidly increase in velocity). That such instability had not been observed, he argued, was evidence against a viscous till rheology.

In order to further constrain the basal mechanics, Engelhardt and Kamb (1998) returned to the UpB camp and deployed a tethered stake apparatus into the surface of the till layer via a borehole. Their aim was to determine what proportion of the ice stream motion might be accommodated by basal sliding, rather than till deformation. Displacement of the tether line indicated that basal sliding *sensu lato* (i.e. potentially including a contribution from shear deformation of the till very close to the ice base) predominated, and may account for as much as 83–100% of the ice stream's motion, with any till deformation restricted to a shear zone 3–25 cm thick at the top of the 9 m thick basal till. If correct (and applicable across the entire ice stream), the implications of this discovery were profound, because the basal boundary conditions and flow law for ice streams would be radically different for basal sliding compared with pervasive till deformation. Engelhardt and Kamb (1998) also pointed out that the erosion, transport and deposition of till would be different under a sliding regime and, although they refrained from any discussion of the geomorphology, it follows that these different mechanisms could produce quite different landscapes beneath ice streams. To further complicate the picture, a similar study from Ice Stream D, also using a tethered stake instrument, showed that most of the ice stream motion (80%) was accommodated by till deformation (Kamb, 2001). Thus, it appeared that the ratio of sliding to till deformation varied from one ice stream to another and also through time, with Kamb (2001) noting how these contrasting results illustrated the 'difficulty of making valid generalisations about the ice stream [flow] mechanism' (p. 174). It also became clear that the flow mechanisms of ice streams located in deep subglacial troughs could be quite different from the Siple Coast ice streams, as earlier work had postulated (McIntyre, 1985). Indeed, several boreholes drilled through Jakobshavns Isbræ, West Greenland, showed that the base of that ice stream is characterised by a layer of temperate ice of substantial thickness that deforms to support a large proportion of the movement of the ice stream. Later seismic investigations by Clarke and Echelmeyer (1996) indicated that much of the basal interface was probably underlain by compacted, non-deforming sediment, confirming the importance of ice deformation (enhanced creep) as the principal flow mechanism.

Suggestions of a near-plastic till rheology under the Siple Coast ice streams were further corroborated by two influential papers by Tulaczyk *et al.* (2000a, b). New triaxial ring-shear tests (and confined uniaxial tests) on the UpB till (Tulaczyk *et al.*, 2000a) confirmed a nearly plastic till rheology, whereby the failure strength of the till was strongly dependent on effective stress, but practically independent of strain and strain rate. Moreover, Tulaczyk *et al.* (2000a) formulated a compressible-coulomb-plastic (CCP) till model that was able to produce the viscous-like vertical distributions of strain in the upper few centimetres of the till layer, that had been suggested based on

the borehole observations (Engelhardt and Kamb, 1998), and which was consistent with textural and compositional properties (Tulaczyk *et al.*, 1998). In the second paper, Tulaczyk *et al.* (2000b) formulated a new analytical ice stream model, termed the 'undrained plastic bed' model, in which the storage of water within the till was an important component. Significantly, this modelling was able to produce two thermo-mechanically controlled states for ice streams, one with a strong bed and low ice velocities, and one with a weak bed and higher velocities. This suggested that ice streams might be susceptible to thermally triggered instabilities, during which small perturbations in the basal thermal energy balance could grow and led to the generation or elimination of the basal conditions required for ice streaming.

A challenge for the plastic bed model was to explain the geomorphology of ice stream beds in terms of the erosion, transport and deposition of sub-ice stream tills. Several workers had argued that a viscously deforming bed was entirely consistent with the production of flutes and drumlins at the ice–bed interface (Alley *et al.*, 1986; Boulton and Hindmarsh, 1987; Rooney *et al.*, 1987a; Hindmarsh, 1998a, b) and till deltas at the grounding line (Alley *et al.*, 1989a). Indeed, Hindmarsh (1998a, b) had demonstrated that, under certain conditions, a viscously-modelled till sheet was unstable and that infinitesimal perturbations in amplitude were able to grow into drumlin-like features that were consistent with the sedimentological features assumed to reflect deforming beds (Hart, 1997). Hindmarsh (1998a, b) clearly acknowledged that, on the small scale, till behaved plastically (Iverson *et al.*, 1997, 1998), but argued that a viscous law represented the simplest model compatible with large-scale observations on ice sheet beds. Hindmarsh argued that the net effect of numerous small-scale failure events (at scales of 0.1 to 1.0 m) is that of viscous behaviour at spatial scales >1 km (see also Hindmarsh, 1997; Fowler, 2003), but this has been questioned more recently based on the stick–slip motion of the lower reaches of Ice Stream B (Tulaczyk, 2006; see also Iverson, 2010).

In contrast, the geomorphological implications of a nearly plastic till rheology were unclear. In order to address this, Tulaczyk *et al.* (2001) proposed a qualitative ploughing model that was consistent with the experimentally-determined Coulomb–plastic rheology and which invoked keels in the base of the ice ploughing through a metres-thick layer of sediment and transporting it downstream. Interestingly, while they acknowledged that there was little direct evidence for the existence of ice keels beneath modern West Antarctic ice streams, they pointed out that earlier geophysical surveys were, at least, consistent with the notion of a 'grooved' till layer (Rooney *et al.*, 1987a; Novick *et al.*, 1994) and that bedforms left behind by palaeo-ice streams strongly supported their assertions (e.g. mega-scale glacial lineations: Clark, 1993; Canals *et al.*, 2000). They presented a series of qualitative arguments to suggest that the keels can be generated by ice passing over a bed that is stronger than the ice (e.g. hard bedrock) and then survive as the ice ploughs through a weaker bed. Like the viscous till model, this ploughing mechanism produces a net till flux down the ice stream, albeit an order of magnitude lower than estimates for pervasive till deformation spanning several metres (see discussion in section 'How is sediment eroded and transported beneath ice streams?').

To summarise, although the detailed geomorphology of ice stream beds remained elusive, attempts had been made to reconcile the two quite different till rheology models with what was known about the conditions at the ice–bed interface under modern ice sheets, and influenced, to some extent, by observations of subglacial bedforms from palaeo-ice sheet beds.

## In search of ice stream sticky spots

Notwithstanding the debates about the rheology of sub-ice stream tills, it had become widely accepted that they offered very little resistance to flow in the areas that had been investigated. Indeed, early calculations of the force balance along ice streams indicated that a significant fraction of the driving stress must be supported by 'side drag' against the slower-moving ice at the lateral margins (Echelmeyer *et al.*, 1994), perhaps approaching as much as 100% on their lower-most ice 'plains' (Bindschadler *et al.*, 1987). However, these values were typically closer to around 50% further upstream (Raymond *et al.*, 2001; their Table 1), which suggested that basal shear stresses must contribute some component of the resistive stresses. Similar results were obtained from a study of the force balance of Rutford Ice Stream (Frolich and Doake, 1988) where it was found that there is a marginal zone in which lateral stresses are high, but that the centre of the ice stream is dominated by basal friction. Moreover, it is unlikely that bed resistance is entirely uniform and Alley (1993) highlighted several lines of evidence from the Siple Coast ice streams that were suggestive of localised areas of higher basal drag, which he referred to as 'sticky spots' (see Table 1 in Alley, 1993). Evidence included ice surface rumples and crevassing (Vornberger and Whillans, 1986), spatially variable till thicknesses (Rooney *et al.*, 1987a), and inversions of basal drag based on ice surface data (MacAyeal, 1992), but their potential geomorphological expression was largely speculative.

argued that large bedrock bumps penetrating a layer of till and protruding into the base of an ice stream were the most likely cause of sticky spots. He argued that for typical ice stream velocities ( $450 \text{ m a}^{-1}$ ), hemispherical bumps of 1 m radius covering <1% of the bed, or 10 m radius covering <2% of the bed, would theoretically be able to support a basal shear stress of 10 kPa. Empirical support for their existence is found from the correspondence between areas of minimum surface velocity and undulations in ice surface morphology that are thought to represent bedrock ridges, e.g. on Ice Stream E (Bindschadler and Scambos, 1991; MacAyeal, 1992; MacAyeal *et al.*, 1995).

Discontinuity in the basal till layer was another possible cause of stickiness identified by Alley (1993), but he pointed out that such discontinuities were unlikely to support very high basal shear stresses because of their tendency to collect a layer of lubricating water from the surrounding till, unless they also possessed abundant large bumps. Nonetheless, there was evidence for their existence beneath modern ice streams. Based on acoustic impedance data, Atre and Bentley (1993) reported laterally varying basal conditions beneath Ice Stream B and C that were inferred to reflect changes in the nature of the sediments composing the bed and/or their physical state (e.g. porosity). Rooney *et al.* (1987a) also noted a 300 m till-free swath (or at least thinner than the seismic resolution of 2 m) from under Ice Stream B, which Alley (1993) suggested could support up to 13% of the basal shear stress in that region. Observations of varying till thicknesses had also been made on another West Antarctic ice stream. Smith (1997a, b) used seismic reflection data to argue that part of the bed of Rutford Ice Stream (running alongside the Ellsworth Mountains and into the Ronne-Filchner Ice Shelf) was characterised by soft, saturated sediments (that may be deforming), whereas other areas were highly consolidated or consisted of poorly lithified sedimentary rock that offered more resistance to ice flow. Later work by Smith (2006) used seismic techniques to detect micro-earthquakes under Rutford Ice Stream and showed that the stiffer areas of the bed that are likely to support basal sliding were much higher friction than areas where thicker regions of

till were inferred to be deforming. Incidentally, more recent work by Smith *et al.* (2015) showed that clusters of microearthquakes most likely represent sticky spots characterised by stiffer/low porosity sediments, but that they accommodate only a small amount of the total basal motion.

Alley (1993) also suggested that raised regions of the ice stream surface could cause moderate increases in shear stress because ice thickness anomalies are around ten times more effective than bed topography in influencing subglacial water flow. Thus, ice surface perturbations can divert water around the underlying region of the bed and cause an increase in till stickiness due to a localised deficit of lubricating water. Alley (1993) suggested that Ice raft 'a' (an elliptical feature 10 km by 5 km and a few metres high, near the mouth of Ice Stream B: cf. Bindschadler *et al.*, 1987), might be an obvious candidate for this type of sticky spot. Elsewhere, under Ice Stream C, it was suggested that areas of well drained ('stickier') till of the order of  $10^2 \text{ m}^2$  generated micro-earthquakes (Anandakrishnan and Bentley, 1993; Anandakrishnan and Alley, 1994). Indeed, the diversion of subglacial water from Ice Stream C and into Ice Stream B has been hypothesised to account for its shutdown (Anandakrishnan and Alley, 1997), although others have invoked a mechanism involving the freeze-on of subglacial water (Price *et al.*, 2001).

Although not considered in Alley's (1993) 'search for ice stream sticky spots', evidence has since emerged that some 'thermal' sticky spots might be caused by the freeze-on of subglacial water. Lliboutry (1987) had recognised that islands of cold-based ice could freeze to high spots in bedrock roughness at scales of  $10^1$  to  $10^3$  m, but he pointed out that this would be unlikely under ice streams where frictional heating would offset any tendency towards freezing. However, modelling showed that ice stream thinning and the advection of cold ice closer to the bed may help promote basal freezing (Payne and Dongelmans, 1997; Christoffersen and Tulaczyk, 2003a, b), and several boreholes from Ice Stream C have encountered a thin layer of frozen till between the basal ice and unfrozen till (Carsey *et al.*, 2002). Evidence of 'freeze-on' may also have been detected under Ice Stream B (see review in Kamb, 2001). Due to thermodynamic feedbacks, it has been hypothesised that basal freeze-on may spread quite rapidly and act to shut-down ice streams (Tulaczyk *et al.*, 2000b; Bougamont *et al.*, 2003a, b).

Thus, soon after the discovery of weak tills beneath ice streams (Blankenship *et al.*, 1986; Engelhardt *et al.*, 1990), a body of evidence had emerged to suggest the presence of localised sticky spots that might explain spatial and temporal variations in ice flow, including their stoppage and reactivation (Anandakrishnan and Alley, 1997; Bougamont *et al.*, 2003a, b). However, while inverse methods, geophysical, and borehole observations provided important evidence of their existence, they lacked the spatial coverage and resolution to confidently identify their cause and distribution and, aside from the inferred presence bedrock bumps and ridges, the geomorphological expression of ice stream sticky spots was largely unknown.

## Summary

Although ice streams were identified in the late 1800s (Rink, 1877), their importance was only recognised as recently as the 1970s, in relation to the potential instability of the WAIS (e.g. Hughes, 1977). This triggered a major growth in their investigation (Figure 1) that began with the pioneering geophysical investigations of the Siple Coast ice streams (Alley *et al.*, 1986; Blankenship *et al.*, 1986), soon followed by direct borehole observations (Engelhardt *et al.*, 1990) and a concerted


effort to understand and model their rapid flow. By the end of the 20th century it had been shown that they flow over a bed of 'soft' sediments and that their motion was accommodated by a combination of deformation within the till layer and/or sliding across its surface, interrupted by the presence of localised sticky spots. Comparisons between different ice streams showed that their basal conditions were highly variable both within and between individual ice streams (Kamb, 2001), and that ice streams that flowed through deep subglacial troughs did not necessarily require a layer of soft deforming sediment (Iken *et al.*, 1993; Clarke and Echelmeyer, 1996). However, our knowledge of the geomorphology of ice stream beds remained very limited. Geophysical investigations (Rooney *et al.*, 1987a) appeared to show buried flutes (or grooves) within till, with several lines of evidence also pointing to the deposition of till deltas at ice stream grounding lines (Alley *et al.*, 1987a; Shabtaie and Bentley, 1987; Alley *et al.*, 1989a). Moreover, these indirect observations were limited in terms of their vertical and horizontal resolution, and were almost exclusively obtained from a very small number of 'pure' ice streams that flowed over relatively flat sedimentary beds and were underlain by several metres of till. Several authors appealed to the notion that observations from palaeo-ice sheet beds would help inform interpretations of subglacial processes beneath ice streams (Boulton, 1986; Murray, 1997), but the study of palaeo-ice stream geomorphology was very much in its infancy and few ice stream beds had been identified with any confidence.

## Identification and Characterisation of the Beds of Palaeo Ice Streams

### In search of Palaeo-ice streams

The first attempt to incorporate the location of ice streams into a palaeo-ice sheet reconstruction appears to be from Denton and Hughes (1981), who depicted numerous ice streams in the former mid-latitude ice sheets of the northern hemisphere, based largely on topographic inference, see Figure 3(a). Their reconstruction clearly benefitted from Hughes' knowledge of West Antarctic ice streams (Hughes, 1977), but Andrews (1982) was more sceptical and, in his review of ice sheet reconstructions, he pointed out that 'it is not known whether or where ice streams existed in the Laurentide Ice Sheet' (p. 25). It is interesting to note, however, that although virtually nothing was known about the geomorphology of ice streams, Luchitta *et al.* (1981) had speculated that ice streams may have carved outflow channels on Mars, based largely on the surface expression of Antarctic ice streams (see Kite and Hindmarsh (2007) for a more recent rendition of this argument).

The importance of incorporating ice streams into palaeo-ice sheet reconstructions was becoming increasingly recognised and Dyke and Prest (1987) included them as convergent flow-lines in their Late Wisconsinan and Holocene reconstruction of the Laurentide Ice Sheet (LIS). There remained, however, a lingering scepticism that ice streams could be identified in the palaeo-record and very few had been identified. Mathews (1991), for example, questioned whether we knew enough about their basal conditions to say anything about the landforms they created, but he drew attention to some features on the continental shelf off Canada that might be considered tracks of former ice streams, including cross-shelf troughs between Vancouver Island and the Queen Charlotte Islands in western Canada (Luternauer and Murray, 1983) and another marine trough extending southeast from the St Lawrence Estuary in



**Figure 3.** (a) Denton and Hughes (1981) hypothesised location of ice streams in the northern hemisphere ice sheets (from Stokes and Clark, 2001). Although largely based on topographic inference and some notion of spatial self-organisation, many of these ice stream locations have since been confirmed in recent inventories derived from detailed mapping of ice stream beds, such as the Laurentide Ice Sheet (Margold *et al.*, 2015a), which is shown in (b), and taken from Stokes (2017). [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]

eastern Canada. He also noted that the large lobate moraines of the southern margin of the LIS (e.g. the Des Moines and Lake Michigan Lobes) may represent another possible type of ice stream track, given their similar size to Antarctic ice streams and their low surface slopes. Mathews (1991) concluded by suggesting that other possible candidates may exist (e.g. in the Canadian Arctic Archipelago) and they merit investigation ‘once the criteria for identifying ice stream tracks become better established’ (p. 267).

Mathews’ (1991) pessimism was, however, misplaced, because Art Dyke and colleagues had already recognised evidence for ice streaming in the central Canadian Arctic (Dyke, 1984; Dyke and Morris, 1988), largely on the basis of carbonate-rich tills dispersed through areas of igneous or metamorphic bedrock. Dyke and Morris (1988) called these ‘Boothia-type’ dispersal trains and described a classic example from Prince of Wales Island that is shown in Figure 4. They noted how the lateral margins of the dispersal train were very abrupt and that a convergent pattern of drumlins feeds into the plume and become more elongated along its central axis, noting that the overall pattern resembled ice streams in

Greenland and Antarctica. Dyke and Morris (1988) also noted a ‘peculiar lateral shear moraine’, a feature unknown from elsewhere’ (p. 86). This feature is at least 68 km long (with minor gaps) and runs parallel to another major drumlin field. Dyke and Morris (1988) speculated that it may have marked a shear zone at the lateral margin of an ice stream.


Thus, around the same time that researchers were striving to glimpse the bed of modern ice streams (Figure 2) in West Antarctica (Blankenship *et al.*, 1986; Rooney *et al.*, 1987a; Engelhardt *et al.*, 1990), Dyke and Morris (1988) were providing the first detailed description of their elusive bed geomorphology (Figure 4) and this paper would go on to have a profound influence on the emerging study of palaeo-ice streams.

### Ice lobes, surging and terrestrial ice streams

Around the same time as Dyke and Morris’ (1988) influential work at the northern margin of the LIS, many of the lobes of the southern margin were attracting attention as possible zones of ice streaming (Dredge and Cowan, 1989) because they had been calculated to possess low surface slopes (Mathews, 1974) that were similar to those in West Antarctica. Hicock and co-workers also reported Boothia-type dispersal trains (Figure 4(a)) and deformation tills associated with drumlins around the Great Lakes region, that were interpreted to reflect ice streaming (Hicock, 1988; Hicock and Dreimanis, 1992). Indeed, Alley *et al.*’s (1986) discovery of a deforming till beneath Ice Stream B led him to conclude that the widespread uniform till sheets of the southern margin of the LIS were also consistent with deposition from deforming subglacial sediment layers beneath relatively short-lived ice streams (Alley, 1991), that had earlier been described as ‘surges’ resulting from the build-up of pressurised subglacial water (Wright, 1973; Clayton *et al.*, 1985). In relation to the Des Moines Lobe, Patterson (1997) argued that the ‘persistence of the lobate margin, the distinctive style of glacial erosion and deposition along the lowland, all led to the interpretation that an ice stream, analogous in size to those in West Antarctica, may have drained a large portion of the central Laurentide Ice Sheet’ (p. 260). Indeed, Patterson (1997) drew heavily on West Antarctic analogues, suggesting that an ice flux similar to that from Ice Stream B ( $30 \text{ km}^3 \text{ a}^{-1}$ ; Bindschadler *et al.*, 1987) could have created the Des Moines lobe in around 1000 years. A further paper by Patterson described specific glacial landform assemblages that characterised several ice streams at the southern Laurentide margin (Patterson, 1998). These included a lowland suite of level-to-streamlined fine-grained till, sometimes associated with highly elongate drumlins (Bluemle *et al.*, 1993) that typically terminated towards the lobe margins, where thrusting of glacial sediment was evident in association with hummocky topography and major moraine systems.

### Giant glacial grooves and mega-scale glacial lineations

The observation of exceptionally long narrow drumlins along some hypothesised ice stream tracks at the southern Laurentide margin (Bluemle *et al.*, 1993) appeared to suggest a link between their formation and rapid ice flow. Giant glacial grooves and other highly elongate landforms had been reported several decades earlier (Smith, 1948; Dean, 1953), but not explicitly linked to ice streaming, which is not surprising given that ice streams had yet to be fully recognised in modern ice sheets (see sections ‘The discovery and definition of ice streams’ and ‘West Antarctic ice streams and “Glaciology’s Grand Unsolved



**Figure 4.** (a) Landsat image (data from Google Earth: Landsat/Copernicus) of a Boothia-type erratic dispersal train on Prince of Wales Island, Canadian Arctic. Dyke and Morris (1988) and Dyke *et al.* (1992) described this classic example and noted how the lateral margins of the dispersal train were very abrupt and coincided with a convergent pattern of drumlins feeding into the plume. (b) Simplified map of the glacial geomorphology from Hättestrand and Kleman (1999), which they redrew from Dyke *et al.* (1992). Reproduced by permission of Elsevier. [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]

Problem"). In central North Dakota, however, Bluemle *et al.* (1993) reported drumlin ridges up to 27 km in length and with typical length to width ratios ranging from 30:1 to 50:1, with a maximum of 240:1. They suggested that the features were formed in basal ice cavities and that sediments were squeezed into the lower-pressure cavity from laterally adjacent regions of the ice stream bed during ice thrusting.

Similar, albeit much larger, patterns of streamlining were also reported by Clark (1993). Using Landsat satellite imagery, he identified a 'hitherto undocumented and much larger form of ice moulded landscape' (p. 1) which comprised streamlined glacial lineations with typical lengths of between 8 and 70 km, widths between 200 and 1300 m and spacings between 300 and 5 km. Clark (1993) termed these features 'mega-scale glacial lineations' (MSGs) and discussed a variety of possible origins, concluding that they probably formed as a result of subglacial deformation and attenuation of sediments around inhomogeneities in till, similar to that earlier proposed for drumlins (Boulton and Hindmarsh, 1987). If correct, he argued that their great length would be a product of the duration of flow and the basal ice velocity, i.e. a 50 km long MSG could form in as little as 50 years under velocities of  $1000 \text{ m a}^{-1}$ , or as much as 5000 years under velocities of  $10 \text{ m a}^{-1}$ . He then proposed that because ice sheet flow-lines are unlikely to have remained stable for long time periods (Boulton and Clark, 1990), MSGs are likely to form under conditions of extremely

rapid flow such as ice streams or surges. This qualitative link between bedform elongation and ice velocity has been further corroborated by more detailed quantitative analysis of drumlins and MSGs on several palaeo-ice stream beds (Hart, 1999; Stokes and Clark, 2002a; Briner, 2007; Stokes *et al.*, 2013a), despite the fact that the precise mechanism(s) of their formation and elongation remains enigmatic (see discussion in section 'How are bedforms created under ice streams?').

### Geomorphological criteria to identify Palaeo-ice streams

Several workers also noted that the flow-sets containing elongate subglacial bedforms were consistent with the presence of palaeo-ice streaming. Hodgson (1994) drew attention to the abrupt lateral margin of a > 100 km long field of highly elongate drumlins on Victoria Island, Canadian Arctic, thought to delimit the position of the lateral shear margin of a palaeo-ice stream (Figure 5), and similar to that proposed by Dyke and Morris (1988) on nearby Prince of Wales Island (Figure 4). Elsewhere, Punkari (1995a, b) postulated the existence of 11 ice streams within a broad marginal zone of the Fennoscandian Ice Sheet (FIS), based largely on satellite imagery that revealed elongate drumlins arranged




**Figure 5.** Landsat image (data from Google Earth: Landsat/Copernicus) of the abrupt lateral shear margin of the M'Clintock Channel Ice Stream on Victoria Island, Canadian Arctic, first identified by Hodgson (1994). [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]

in fan-shaped flow patterns that contrasted with glaciofluvial landform assemblages in the inter-lobate zones (Punkari, 1997). Similar fans were described by Kleman and Borgström (1996) who termed them 'surge fans' and noted their distinctive bottle-neck flow patterns (convergent and then divergent) of glacial lineations that probably form rapidly and during the decay stages of an ice sheet. Kleman and Borgström (1996) identified the Dubawnt Lake surge fan in central Canada as a

'type landscape, which has subsequently been shown to represent a major, but short-lived, ice stream track during final deglaciation of the Keewatin sector of the LIS (Stokes and Clark, 2003).

In terms of the geomorphology of ice streams, therefore, significant progress was made during the 1990s in terms of identifying evidence of ice streaming among the glacial geomorphological record of palaeo-ice sheet beds. These studies suggested that ice streaming should leave behind sedimentological evidence of fast ice flow in the form of heavily deformed tills and distinctive erratic dispersal trains that often depicted convergent flow-patterns (Dyke and Morris, 1988; Hicock, 1988; Alley, 1991; Patterson, 1997, 1998). Many of these flow-sets, or fans (cf. Kleman and Borgström, 1996), also contained highly elongate glacial lineations (MSGs), which were postulated to reflect rapid ice velocities (Clark, 1993); and some were characterised by abrupt lateral margins (Hodgson, 1994) and lateral shear margin moraines (Dyke and Morris, 1988; Stokes and Clark, 2002a). Taken together, these were argued to represent the key 'geomorphological criteria' for identifying palaeo-ice streams, which Stokes and Clark (1999) encapsulated in a series of landsystems models (shown in Figure 6) depending on whether the ice stream terminated in water or on land, and whether the glacial lineations were formed rapidly and synchronously or time-transgressively during ice margin retreat. Stokes and Clark (1999) hoped that these observational templates would aid the identification of ice streams in palaeo-ice sheets. Indeed, in the first review of the evidence of palaeo-ice streams in 2001, it was clear that only a very small proportion of palaeo-ice streams had been identified with any confidence (Stokes and Clark, 2001). In part, this may have been because all modern ice streams are marine-terminating and unlike, for example, surge-type glaciers, it was difficult to observe the geomorphology in their recently deglaciated foregrounds and identify modern analogues (Evans *et al.*, 1999). As a result, Stokes and



**Figure 6.** Idealised landsystems models of the geomorphology of palaeo-ice stream tracks from Stokes and Clark (1999) depending on whether the ice stream was marine-terminating (a and b) or terrestrially-terminating (c and d) and whether the imprint was formed near-synchronously prior to ice stream shut-down (a and c) or time-transgressively during ice stream retreat (b and d). These imprints were designed to help workers identify palaeo-ice stream tracks on the beds of palaeo-ice sheets. Reproduced from Benn and Evans (2010) with permission from Hodder Education, London, UK. [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]


Clark (2001) suggested that geophysical observations of the submarine glacial geomorphology recently exposed in front of modern-day ice streams (Shipp *et al.*, 1999) held huge potential to link the terrestrial record of mid-latitude palaeo-ice streams with their contemporary counterparts in West Antarctica.

### Offshore records of palaeo-ice streams based on marine geophysics


The early work identifying palaeo-ice stream footprints was based largely on terrestrial glacial geology and was considerably aided by advances in satellite remote sensing in the early 1990s (see examples in Figures 4 and 5). In a similar manner, advances in marine geophysical techniques during the late 1990s opened up new research aimed at identifying the geomorphology of palaeo-ice streams offshore (Ó Cofaigh, 2012). Early work using seismic and side-scan sonar focused on glacially influenced continental slopes in the Polar North Atlantic (e.g. under the auspices of the 'Polar North Atlantic Margins' (PONAM) programme). Key contributions related to ice stream geomorphology included the discovery of large sedimentary fans at the mouth of cross-shelf troughs (termed 'Trough Mouth Fans': Vorren and Laberg, 1997), whose volume and architecture were interpreted to reflect rapid sedimentation by ice streams (Dowdeswell *et al.*, 1996; Hooke and Elverhøi, 1996; King *et al.*, 1996; Elverhøi *et al.*, 1997; Vorren and Laberg, 1997; Vorren *et al.*, 1998; Batchelor and Dowdeswell, 2014). In many cases, rapid sedimentation was interpreted to reflect transport via a deformable bed that was subsequently deposited at the ice stream grounding line and then remobilised down-slope (see Figure 7), consistent with the observations and modelling of modern ice streams (Alley *et al.*, 1986, 1987a, b, 1989a, b; see section 'Ice stream geophysics: a paradigm shift in glaciology?'). One of the key advantages of these marine records is that the architecture of the depocentres could be used to bracket the age of sediment deposition and, hence, rates of sediment flux (Dowdeswell *et al.*, 2004; Nygård *et al.*, 2007). These estimates typically span an order of magnitude from  $100 \text{ m}^3 \text{ a}^{-1}$  (Alley *et al.*, 1989a) up to  $1000 \text{ m}^3 \text{ a}^{-1}$  per metre width of ice stream terminus (Dowdeswell *et al.*, 2004), although far more extreme rates have also reported, e.g.  $8000 \text{ m}^3 \text{ a}^{-1}$ , which are comparable with some of the world's largest rivers (Nygård *et al.*, 2007).

A further advantage is that marine-based seismic data can provide spatially extensive information of sub-surface sediment characteristics of the ice stream geomorphology, which can often be used in conjunction with geotechnical analysis of sediment cores. In this regard, several studies have shown that the uppermost layer of the sub-surface of the former ice stream track is typically composed of an acoustically transparent till that is generally under-consolidated compared with the material beneath, see Figure 8 (Dowdeswell *et al.*, 2004; Ó Cofaigh *et al.*, 2005, 2007; Nygård *et al.*, 2007). This is often interpreted as a low shear strength ('soft') deformation till that facilitated the rapid basal motion of the ice stream (Dowdeswell *et al.*, 2004; Ó Cofaigh *et al.*, 2005, 2007) and which is subsequently deposited at the terminus (Figure 7). Marine geophysical surveys of palaeo-ice stream beds indicates that the thickness of this till layer typically averages  $\sim 5\text{--}10 \text{ m}$  (Dowdeswell *et al.*, 2004; Ó Cofaigh *et al.*, 2005, 2007; Livingstone *et al.*, 2016a), which is similar to the inferences from the early geophysical explorations of modern West Antarctic ice streams (see section 'Ice stream geophysics: a paradigm shift in glaciology?'). Detailed analyses of these metres-thick 'soft' tills, however, has shown that they are typically characterised by a range of structures indicative of subglacial shear and that such shearing tends to be concentrated into thin (0.1–0.5 m thick) zones, indicating that deformation was unlikely to be pervasive throughout the entire thickness and was more consistent with localised near-plastic rheology (Ó Cofaigh *et al.*, 2005). These datasets also reveal that MSGs are typically formed in this till layer (Figure 8), and recent work suggests that MSGs are clustered more closely and reach greater lengths in areas of relatively thicker till (6–12 m), compared with areas where the soft till is thinner or where the till is stiffer (Livingstone *et al.*, 2016a).

The increasing resolution of swath bathymetry data from the late 1990s also allowed workers to map the large-scale distribution of landform-sediment associations on continental shelves surrounding present-day ice sheets in Antarctica (Wellner *et al.*, 2001) and Greenland (Evans *et al.*, 2009) and those formerly occupied by mid-latitude palaeo-ice sheets in North America (Shaw *et al.*, 2006) and NW Europe (Ottesen *et al.*, 2005, 2008). This allowed numerous palaeo-ice streams to be identified based on the location of cross-shelf troughs separated by zones of slower-flow on the adjacent banks (see recent inventories by Ottesen *et al.*, 2005; Livingstone *et al.*, 2012; Batchelor and Dowdeswell, 2014; Margold *et al.*, 2015a).




**Figure 7.** Schematic model showing sedimentary processes and the development of a till delta at grounding line of a marine-terminating ice stream at the edge of the continental shelf (e.g. during the Last Glacial Maximum) (redrawn from Laberg and Vorren, 1995).



**Figure 8.** TOPAS sub-bottom profiler records from outer Marguerite Trough palaeo-ice stream, Antarctic Peninsula (from Ó Cofaigh *et al.*, 2005) showing mega-scale glacial lineations formed in an acoustically-transparent, low shear strength till averaging around 5–10 m thick. Reproduced by permission of Elsevier.

Collectively, these studies also indicated that palaeo-ice sheets likely had very similar velocity patterns to modern ice sheets (Margold *et al.*, 2015b). Moreover, these studies revealed that the geomorphology within the cross-shelf troughs was similar to that observed in terrestrial settings (see section ‘Geomorphological criteria to identify palaeo-ice streams’), with MSGLs typically observed on the mid- to outer-cross-shelf

troughs and sometime in association with lateral shear margin moraines, see Figure 9 (Wellner *et al.*, 2001; Ó Cofaigh *et al.*, 2002; Ottesen *et al.*, 2005, 2008). This helped to strengthen the association between specific glacial landforms (e.g. MSGLs, ice stream shear margin moraines) and ice streaming, particularly where they are observed in troughs that lie immediately in front of modern-ice streams (Shipp *et al.*, 1999; Wellner



**Figure 9.** (a) Swath bathymetry of the Malangsdjupet palaeo-ice stream bed in northern Norway (from Ottesen *et al.*, 2008) showing the typical geomorphology of a marine-terminating ice stream with rough bedrock in the onset zone merging into drumlins and mega-scale glacial lineations in the deepest part of the cross-shelf trough. Also note the location of a grounding zone wedge (cross section from seismic data shown in (b)) and lateral shear margin moraines (black arrows). Reproduced by permission of Elsevier. [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]




Larter *et al.*, 2009). Graham *et al.* (2009) also recognised clear spatial variability in the landform signature, such that bedforms of quite different size and shape can occur in close proximity, which might be related to local changes in basal conditions (e.g. sticky spots), particularly in the inner to mid-shelf areas.

Marine geophysical records of formerly marine-based palaeo-ice streams have also identified landforms that lie transverse to ice flow and which often occur superimposed on the flow-parallel bedforms. The largest and most prominent of these are grounding zone wedges (GZWs), which are asymmetric sediment depocentres (with steeper ice-distal slopes) typically several kilometres long (down-flow) and ranging in thickness from 10 to 100 m (Figure 9: Dowdeswell and Fugelli, 2012; Batchelor and Dowdeswell, 2015). They are typically composed of subglacial till and often contain seaward dipping reflectors which indicate progradation of sediment as it is deposited at the grounding line, as predicted by Alley *et al.* (1986), who suggested ‘till deltas’ would be a consequence of their deforming bed model (section ‘Ice stream geophysics: a paradigm shift in glaciology?’, Figure 7). The formation of GZWs is, therefore, thought to require high rates of sediment transport to a grounding line position that is relatively stable for 10s to 100s years (Graham *et al.*, 2010; Dowdeswell and Fugelli, 2012; Batchelor and Dowdeswell, 2015; Livingstone *et al.*, 2016a). For example, radiocarbon dates that bracket

the age of GZW formation on the outer shelf of Marguerite Trough require sediment fluxes  $>1000\text{ m}^3\text{ a}^{-1}$  (per meter width of the grounding line) (Livingstone *et al.*, 2016a). Because their formation also requires a stable grounding line position, at least for a few decades, it has been noted that they tend to form in locations where the geometry of the ice stream’s terminus would have been constricted by a narrower or shallower topographic setting (Graham *et al.*, 2010; Jamieson *et al.*, 2012; Batchelor and Dowdeswell, 2015). Although progradation of a sediment via a deforming layer is often invoked to account for GZW formation, Dowdeswell and Fugelli (2012) noted that channel-like incisions are associated with some of these features and may suggest that delivery of sediment via pressurised subglacial meltwater, perhaps associated with drainage of subglacial lakes (see section ‘Direct observations of subglacial hydrology and till deltas’).

Narrower recessional moraines have also been identified superimposed on marine-based palaeo-ice stream beds (Shipp *et al.*, 2002; Graham *et al.*, 2010). These are typically  $<15\text{ m}$  thick and  $300\text{ m}$  wide and possess sediment volumes that are an order of magnitude less than GZWs (Batchelor and Dowdeswell, 2015). They occur in assemblages containing 10s to 100s of parallel to sub-parallel ridges and are thought to form by ice pushing of sediment, including folding, faulting and thrusting, during minor ice sheet re-advances within



**Figure 11.** Idealised landsystems models of the geomorphology of palaeo-ice stream tracks that have undergone rapid retreat (top), episodic retreat (middle) and slow retreat (bottom) (from Ó Cofaigh *et al.*, 2008). Reproduced by permission of John Wiley and Sons Ltd.

overall retreat (Shipp *et al.*, 2002; Ottesen and Dowdeswell, 2006, 2009). The significance of both GZWs and transverse ridges is that they record the style of retreat of marine-based ice streams, see Figure 11. The absence of these features is often taken to infer rapid retreat of the ice stream, with sporadic occurrences of GZWs taken to infer episodic retreat punctuated by still-stands, and sequences of minor recessional moraines taken to infer relatively slow retreat of a grounded ice margin (Figure 11) (Dowdeswell *et al.*, 2008; Ó Cofaigh *et al.*, 2008).

More recently, a smaller-scale series of transverse ridges (termed 'corrugations': Graham *et al.*, 2013) have been identified in high resolution swath bathymetry on some palaeo-ice stream beds (Jakobsson *et al.*, 2011; Graham *et al.*, 2013). These ridges are just a few metres high and their crests are separated by ~60 to 200 m, with a spacing observed to decrease seaward (Jakobsson *et al.*, 2011). They are thought to be formed by the periodic and tidally-modulated grounding of large icebergs (cf. Lien *et al.*, 1989) within an ice mélange, and have been hypothesised to result from rapid disintegration of an ice shelf (Jakobsson *et al.*, 2011). However, Graham *et al.* (2013) proposed a more generic origin that involves the forward motion of a ploughing ice keel that might occur beneath an ice shelf, or in the vicinity of the grounding line, or behind newly calved icebergs at the grounding line.


### In search of Palaeo-ice stream sticky spots

As noted in section 'In search of ice stream sticky spots', several studies had confirmed the existence of sticky spots beneath active ice streams (Anandakrishnan and Alley, 1994; MacAyeal *et al.*, 1995; Price *et al.*, 2002; Joughin *et al.*, 2004), but their geomorphological expression was largely unknown. This point was emphasised by MacAyeal *et al.* (1995: p. 247) who stated that 'if sticky spots exist and, if their influence on ice stream flow is important, the characterisation of sticky spots (emphasis added) may prove essential to understanding how ice sheets evolve with time'. As a result, several workers recognised that the geomorphology on the beds of palaeo-ice streams offered huge potential to characterise their nature and distribution (see review in Stokes *et al.*, 2007). Thus, it was argued that where subglacial bedforms (drumlins and MSGs) were shorter and/or deviated around areas of exposed bedrock or topographic highs on the ice stream bed, that these patterns represented sticky spots (Hodgson, 1994; Clark and Stokes, 2001; Stokes *et al.*, 2007). It was also noted that these areas were particularly common in the onset zones of marine-based ice streams (e.g. Figures 9, 10) where ice flow was likely accelerating, but had not reached typical streaming velocities (Canals *et al.*, 2000; Wellner *et al.*, 2001; Ó Cofaigh *et al.*, 2002; Shipp *et al.*, 2002; Graham *et al.*, 2009). However, Clark and Stokes (2001) noted a more unusual downstream decrease in bedform elongation on the M'Clintock Channel Ice Stream bed in the Canadian Arctic Archipelago (Figure 5), whereby MSGs transitioned into shorter drumlins that were located among areas of exposed bedrock. These bedrock-dominated areas were interpreted as till-free sticky spots (cf. Stokes *et al.*, 2007) and Clark and Stokes (2001) hypothesised that 'frictional shut-down' of the ice stream occurred when the soft lubricating sediments at the bed became non-existent. The small incorporation of erratic material in till samples from the ice stream bed (~4%: see Hodgson, 1993) also indicated that sediment supply from upstream was minimal, highlighting the importance of till continuity for the operation of some ice streams (cf. Alley, 2000).

Variations in the routing and pressure of subglacial meltwater were also becoming increasingly linked to variations in glacier

bed stickiness and the concept of a 'mosaic' of deforming and stable spots was becoming widely accepted (Boulton *et al.*, 2001; Piotrowski *et al.*, 2004). Such variations had already been invoked to explain the shut-down of Ice Stream C, e.g. the re-routing of subglacial meltwater and/or its freeze-on to the base of the ice (Anandakrishnan and Alley, 1994; Bougamont *et al.*, 2003a). The sedimentological evidence of the deforming bed mosaic was described in Piotrowski *et al.* (2004) and evidence of 'stick-slip' behavior had been described in tills from several palaeo-ice stream beds (Hicock and Dreimanis, 1992; Knight, 2002; Lian *et al.*, 2003). The geomorphological expression of this type of sticky spot, however, was largely unknown, but several workers noted the rather unusual occurrence of transverse features, known as ribbed moraine, superimposed on highly elongate bedforms that were the classic signature of ice stream flow (Figure 4: Dyke and Morris, 1988; Dyke *et al.*, 1992)

Ribbed moraines form transverse to ice flow and generally occur in fields of closely spaced ridges of the order of 10 m high, 100 m wide and 1000 m long (Hättestrand and Kleman, 1999; Dunlop and Clark, 2006). Stokes *et al.* (2007) noted that they are associated with palaeo-ice stream beds in two situations: (i) exclusively in the onset zone, merging into drumlins and MSGs that exist downstream (Figure 4: Dyke and Morris, 1988; Dyke *et al.*, 1992; De Angelis and Kleman, 2008); and (ii) in discrete patches within the main ice stream trunk, clearly superimposed on top of MSG and drumlins, see Figure 12 (Stokes and Clark, 2003; Stokes *et al.*, 2006). Their occurrence on ice stream beds is unusual because they are typically found in the interior regions of ice sheets (Aylsworth and Shilts, 1989; Hättestrand, 1997; Kleman and Hättestrand, 1999). Thus,



**Figure 12.** Landsat image (data from Google Earth: Landsat / Copernicus) of ribbed moraines superimposed on mega-scale glacial lineations in the main trunk of the Dubawnt Lake Ice Stream bed. It has been hypothesised that the ribbed moraines are a manifestation of sticky spots during ice stream shut-down (Stokes and Clark, 2003). [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]

Stokes *et al.* (2007) argued that their appearance likely indicates an abrupt change in ice dynamics and most likely a switch to 'slow flow' that is related to the development of sticky spots.

In ice stream onset zones, ribbed moraines have been reported to develop at the transition between the cold-based catchment and warm-based ice stream (Dyke and Morris, 1988; Dyke *et al.*, 1992; Hättestrand and Kleman, 1999; De Angelis and Kleman, 2008). Dyke and Morris (1988) were the first to document ribbed moraines in the onset zone of a palaeo-ice stream on Prince of Wales Island in the Canadian Arctic Archipelago (Figure 4) and they invoked oscillations of the basal thermal regime between cold and warm-based ice that led to acceleration and deceleration of ice velocity and attendant infolding and stacking of debris. De Angelis and Kleman (2008) also noted that the geomorphological imprint of four ice stream onset zones they studied in the NE Laurentide Ice Sheet (including the one studied by Dyke and Morris, 1988) matched the typical characteristics of partially frozen beds, e.g. palimpsest fluvial landscapes or older glacial landscapes juxtaposed with areas with fully developed glacial lineations and often associated with ribbed moraines (Kleman and Borgström, 1996).

In contrast, there are a few cases where ribbed moraines occur in the main trunk of the ice stream. In the case of the Dubawnt lake Ice Stream bed, central Canada (see Figure 12), Stokes and Clark (2003) reported their patchy appearance superimposed on MSGs, and suggested that they formed through glaciotectionic shearing and stacking of debris under compressional ice flow that resulted from a loss of water in the till layer, perhaps related to basal freeze-on (cf. Stokes *et al.*, 2008). More recently, Trommelen and Ross (2014) have also linked ribbed moraines to sticky patches with the subglacial bed mosaic of northern Manitoba. They argued that 'pristine' ribbed moraines were preserved under cold-based ice, but that other patches of ribbed moraine were partially to fully streamlined during a transition to warm-based ice during deglaciation. Interestingly, their detailed study of till composition and dispersal under the Hayes Lobe (terrestrial ice stream) in NE Manitoba (Trommelen *et al.*, 2014) revealed the presence of sticky spots that were identifiable as islands of 0.2 to 4 km wide patches of till dominated by a local composition, but surrounded by a regional farther-travelled till sheet. The cause of these inferred sticky spots is intriguing because they occur in an area of abundant warm-based streamlined bedforms, but they are possibly related to variations in subglacial water pressure (Trommelen *et al.*, 2014).

Most recently, Winsborrow *et al.* (2016) noted the presence of glaciotectionic rafts and hill-hole pairs on an ice stream bed associated with the former Barents Sea Ice Sheet and argued that they represent the geomorphological expression of sticky spots. In this location on the continental shelf, however, they noted that the sticky spots coincided with sub-surface shallow gas accumulations and that the formation of gas hydrates could have desiccated, stiffened, and thereby strengthened the subglacial sediments, which led to the development of a sticky spot. On the basis of these observations, they suggested that a further cause of sticky spots is likely related to pore-water piracy and sediment stiffening due to subglacial gas-hydrate accumulation, and that such a process could be more prevalent under extant ice streams than hitherto recognised.

## Summary

Around the same time that ice streams were gaining prominence in glaciology in the 1980s, glacial geomorphologists were attempting to identify their 'footprints' on the beds of

palaeo-ice sheets. Denton and Hughes (1981) were the first to fully recognise the importance of ice streams to understanding past ice sheet dynamics, but their reconstruction of ice stream locations in Northern Hemisphere ice sheets (Figure 3(a)) was initially met with scepticism, which is perhaps understandable given that there were no criteria to objectively identify ice stream footprints in the palaeo record, and very few had been identified based on empirical evidence (see section 'Ice stream geophysics: a paradigm shift in glaciology?'). During the late 1980s, however, pioneering work on erratic dispersal trains (e.g. Figure 4: Dyke and Morris, 1988) was soon followed by the recognition that ice streams created a distinct geomorphology that included convergent bedform patterns, highly elongated glacial lineations (MSGs), ice stream shear margin moraines, and major sedimentary depocentres (till deltas, grounding zone wedges, trough mouth fans). Collectively, these geomorphological criteria (cf. Stokes and Clark, 1999) formed a series of landsystem templates (e.g. Figures 6, 10 and 11) that enabled their objective identification on palaeo-ice sheet beds. The increased use of satellite remote sensing and marine geophysical techniques led to the identification of hundreds of palaeo-ice streams, both onshore and offshore, such that some inventories are probably close to complete (Ottesen *et al.*, 2005, 2008; Livingstone *et al.*, 2012; Batchelor and Dowdeswell, 2014; Margold *et al.*, 2015a, b; Dowdeswell *et al.*, 2016).

By the early 2000s, we knew what a former ice stream bed looked like in terms of its geomorphology, including the key diagnostic landforms of rapid ice flow – mega-scale glacial lineations (Clark, 1993) – and new insights regarding how sticky spots might be manifest in the geomorphological record. These landforms have been used to reconstruct where and when ice streams operated (Ottesen *et al.*, 2005; Stokes *et al.*, 2016a) and as both input and constraint data for numerical modelling of individual ice streams (Jenson *et al.*, 1995; Jamieson *et al.*, 2012) and ice sheets (Stokes and Tarasov, 2010). These observations provided a clear geomorphological framework for those studying the beds of modern ice streams using geophysics (section 'Early Exploration of the Beds of Active Ice Streams') and led to increased cross-pollination of ideas between the two disciplines. However, palaeo-ice stream beds are a time-integrated and often fragmentary record of ice stream operation that these data could say little about the time-scale over which processes were operating. Indeed, palaeo-ice stream research was in its infancy with respect to the time-scales over which the geomorphology was created and evolved and under what ice dynamical conditions (ice velocity, thickness, etc.). Smith *et al.* (2007: p.127) also noted that 'the interpretation of former ice dynamics from the subglacial bedforms left behind following deglaciation is also limited by a lack of observed analogous features beneath modern glaciers'. This would require a renewed effort to observe the geomorphology and basal conditions on modern ice stream beds using geophysical techniques with much higher resolution.

## Geophysical Investigation of the Geomorphology beneath Modern Ice Streams


### Discovery of subglacial bedforms beneath modern ice streams

Soon after the early work on the 'pure' Siple Coast ice streams in West Antarctica (section 'The discovery and definition of ice streams'), efforts were underway to investigate the basal conditions beneath Rutford Ice Stream, also in West

Antarctica (King *et al.*, 2004; Smith, 1997a, b; Smith *et al.*, 2007). Unlike the Siple Coast ice streams, it flows through a deep asymmetric bedrock trough, bounded on one side by the Ellsworth Mountains, and with ice thicknesses ranging from ~2000–3200 m and ice velocities 300–400 m a<sup>-1</sup> (Smith, 1997a, b; King *et al.*, 2007). As such, its characteristics are more typical of the majority of extant ice streams in Greenland and Antarctica that are, to a large extent, controlled by underlying topography. Building on the earlier work on this ice stream (Stephenson and Doake, 1982; Doake *et al.*, 1987), Smith (1997a, b) used seismic reflection profiles from transects perpendicular to the ice stream bed to show that the characteristics of the bed material varied both along and across the ice stream. The majority of the bed was interpreted as dilatant water-saturated sediments undergoing pervasive deformation, but other areas were interpreted to be non-deforming and perhaps accommodating basal sliding (Smith, 1997a). Smith (1997b) also noted that the proportion of the ice stream width over which bed deformation occurred appeared to increase downstream over a distance of ~50 km (see also Doake *et al.*, 2001). Localised areas of very high basal shear stress (sticky spots) were also inferred (see later work by Smith, 2006), although Smith (1997a) noted little correspondence between these areas of the ice stream bed and surface velocity data or satellite imagery of the ice stream surface. Of significance to ice stream geomorphology was that, although the profile data were only collected in two dimensions, Smith (1997a) tentatively interpreted a 'bump' approximately 400 m wide and 50 m high as a drumlin composed of soft deforming sediments sitting on a more rigid substrate. This appears to represent the first direct evidence of a subglacial bedform beneath an active ice stream.

Later work by Smith *et al.* (2007) collected seismic reflection data from the same location as those presented in Smith (1997a). In addition to the data collected from the original lines in 1991, Smith *et al.* (2007) presented data from 1997 and 2004, when the ice stream flow had not changed significantly. The repeat measurements enabled them to detect changes in the bed topography (geomorphology) and associated changes in subglacial hydrology (from acoustic impedance) over a 13 year period, shown in Figure 13. Between 1991 and 1997, they measured erosion of sediment averaging 1 m a<sup>-1</sup> across a 500 m wide area, which Smith *et al.* (2007) noted was much higher than previous estimates of subglacial erosion in other subglacial environments and under the Siple Coast ice streams (<0.1 to 100 mm a<sup>-1</sup>: Alley *et al.*, 1987b, 1989a; Humphrey and Raymond, 1994; Hallet *et al.*, 1996; Alley *et al.*, 2003), although more comparable with recent work from Pine Island Glacier, which estimates similarly high rates of 0.6 ± 0.3 m a<sup>-1</sup> (Smith *et al.*, 2012). They ruled out the possibility that the erosion was due to subglacial meltwater because any free water at the ice–bed interface would be expected to occur elsewhere in the deeper parts of the ice stream's trough, but they were unable to pinpoint the exact mechanism through which the sediment was removed. The second comparison between the lines acquired in 1997 and 2004 indicated that the erosion across this area had ceased and that a mound of material 10 m high and 100 m wide appeared (Figure 13). An additional flow-parallel line from 2004 also suggested that the mound extended at least 1 km upstream. Smith *et al.* (2007) noted that the dimensions and sedimentary characteristics of the mound were typical of a drumlin, thereby confirming the earlier supposition from Smith (1997a).

Ascertaining the mechanism of drumlin formation was more difficult, but Smith *et al.* (2007) considered two possibilities. The first invoked an advecting groove in the base of the ice that was infilled by soft deforming sediment, similar to that



**Figure 13.** Bed topography and basal conditions at the base of Rutford Ice Stream, West Antarctica (reproduced from Smith *et al.*, 2007, courtesy of the Geological Society of America). (a) Bed topography compared between 1991 and 1997. (b) bed topography between 1997 and 2004 and from an upstream line in 1997. (c) Inferred basal conditions from acoustic impedance data. Ice flow is into the page and error bars are given to the left of each of the lines in (a) and (b).

suggested for flutes (cf. Boulton, 1976). The second invoked a rheological instability in the bed material that had been modelled by Hindmarsh (1998a, b). The former mechanism was favoured on the basis that the feature was longitudinally continuous as well as relatively invariant with time (in both geometry and properties). Notwithstanding these uncertainties, the key conclusion was that an ice stream can reorganise its bed rapidly over just a few years or less, and the direct observation of the formation of a subglacial bedform had broken new ground in the study of ice stream geomorphology.

Around 160 km further upstream, in the onset zone of Rutford Ice Stream, King *et al.* (2007) presented both seismic and radar data to image the bed in a region where the ice flow velocity accelerated from 72 to >200 m a<sup>-1</sup>. In the slower-moving part they observed a transverse moraine 2 km wide and 1.5 km long, which they interpreted to be composed of unconsolidated sediment undergoing active deformation. King *et al.* (2007) suggested that this feature might be analogous to a drumlinised ribbed moraine (Dunlop and Clark, 2006; Trommelen *et al.*, 2014). About 30 km downstream, where velocities exceeded 95 m a<sup>-1</sup>, they observed several 'drumlins of classical form' with elongation ratios between 1:1.5 and 1:4 (King *et al.*, 2007: p. 665). Based on the composition of the drumlins and their conformity with the ice base, they interpreted the drumlins as active depositional features that were similar to the feature observed by Smith *et al.* (2007), and were consistent with erosion and deposition within a mobile deforming bed (Boulton and Hindmarsh, 1987). In addition to the confirmation that bedforms were being created under the ice stream, King *et al.*'s (2007) study was significant because it was the first to demonstrate a clear link between ice velocity and bedform elongation that had only been hypothesised from work on palaeo-ice stream beds (see section 'Giant glacial grooves and mega-scale glacial lineations'), i.e. the drumlinised ribbed moraine were found under ice velocities of ~72 m a<sup>-1</sup>, whereas the more elongate drumlins were observed beneath ice flowing at ~125 m a<sup>-1</sup> (see also Smith and Murray, 2009). King *et al.* (2007) also noted that the

drumlin further downstream (observed by Smith *et al.*, 2007) was lower (10 m vs 30–50 m) and narrower (100 m vs 200–500 m) than the drumlins they had imaged in the onset zone.

### Formation of MSGs beneath a West Antarctic ice stream


Interestingly, Smith *et al.* (2007) had ended their paper documenting drumlin formation under Rutford Ice Stream by suggesting that future work might ascertain ‘whether or not the drumlin evolves into a mega-flute’ (p. 130). This point was, perhaps, a tacit acknowledgement that subglacial bedforms under the fast-flowing trunk of the ice stream would be expected to be much longer than drumlins and more similar to the MSGs (cf. Clark, 1993) that had been observed on palaeo-ice stream beds (section ‘Giant glacial grooves and mega-scale glacial lineations’). Ascertaining the presence of MSGs, however, would require data from a much larger area and Smith and Murray (2009) were able to do this by combining a number of previously-collected seismic lines (reported in Smith, 1997a, b) distributed over an area of 140 km<sup>2</sup>. Interpolation of these datasets allowed them to construct a schematic cross-section of the ice stream bed, shown in Figure 14, which indicated dilatant deforming till in two troughs that run either side of a central high. The central high was characterised by a mosaic of deforming and stiffer till (cf. Piotrowski *et al.*, 2004; see section ‘In search of palaeo-ice stream sticky spots’), and they noted a local downstream transition from deforming to stiffer till, where they inferred basal sliding was more important. They confirmed the location of previous mounds and drumlins (see Smith, 1997a, b; Smith *et al.*, 2007) and noted how some of the deforming material associated with these bedforms extended downstream over the stiffer till. Significantly, they also noted that the ‘bump’, previously interpreted as a drumlin (Smith 1997a, b), continued downstream for at least 17 km and, as a result, they suggested that it should be referred to as a MSG (Clark, 1993).

Subsequently, King *et al.* (2009) acquired new high resolution radar data from a large area of the ice stream bed, including the area where previous seismic lines had been acquired (Smith *et al.*, 1997a, b; Smith and Murray, 2009). Combining the new radar data with the previously-collected seismic data clearly revealed an assemblage of MSGs


beneath ice flowing at around 375 m a<sup>-1</sup>. This assemblage, shown in Figure 15, was characterised by a pattern of ridges and troughs with wavelengths transverse to ice flow of 300 to 1000 m and with peak-to-trough amplitudes ranging from 5 to 90 m, with a mean of 10 m. The longest ridges extended for >18 km and had elongation ratios ranging from 15:1 to >35:1. Indeed, the MSGs observed under Rutford Ice Stream were largely indistinguishable from those observed on palaeo-ice stream beds (Figure 15(b)) and this study provided the first conclusive evidence that MSGs are diagnostic of ice stream flow, which had been postulated well over a decade earlier (Clark, 1993). The King *et al.* (2009) ‘bed-map’ has recently been extended in the downstream direction to cover an area 18 × 40 km and is now available as a dataset that comprises both ice surface, ice thickness and bed elevation data (see King *et al.*, 2016)

As in previous work (Smith, 1997a, b; King *et al.*, 2007; Smith and Murray, 2009), the MSGs were inferred to be developed in areas of dilatant deforming till that formed part of a dynamic sedimentary system that underwent both erosion and deposition on decadal time-scales. King *et al.* (2009) also confirmed that the basal sliding zones, where the till is stiffer and sliding is inferred to be taking place across the surface of the sediment, coincided with areas of the bed that had a more subdued topography and with fewer and poorly streamlined bedforms. The binary distribution of acoustic impedance data were also deemed to be compatible with observations from marine geophysical observations of MSGs formed in a dilatant ‘soft’ till overlying a more consolidated and stiffer till (Figure 8) (Dowdeswell *et al.*, 2004; Ó Cofaigh *et al.*, 2005, 2007). King *et al.* (2009) also noted that none of the existing theories of MSGL formation can readily explain the observations, but they favoured a till instability mechanism (cf. Hindmarsh, 1998a, b) based on the close association between MSGs and the areas of dilatant deforming sediment.

Advances in geophysics were also allowing other ice stream beds to be imaged at higher resolution. Jezek *et al.* (2011) applied radar tomography methods to very high frequency airborne synthetic aperture radar data to measure ice thickness on Jakobshavn Isbræ in West Greenland and construct a 3-dimensional map of the bed topography over an area 5 × 20 km. They were able to detect, for the first time beneath the Greenland Ice Sheet, assemblages of elongate ridge-groove landforms oriented in the direction of the ice flow. These had wavelengths of 150 to 500 m, amplitudes of 10 to 30 m and



**Figure 14.** Schematic cross-section and reconstruction of the geomorphology at the bed of Rutford Ice Stream, West Antarctica (from Smith and Murray, 2009). The line labelled C1 is the same profile as shown in Figure 13. Reproduced by permission of Elsevier.



**Figure 15.** (a) Three dimensional image of the bed of Rutford Ice Stream (from King *et al.*, 2009) looking downstream (see also Figure 14), with the colour shading showing the difference between the short-wavelength topography and a long wave-length trend surface (example radar profile shown top left). (b) Comparison between the mega-scale glacial lineations on the Rutford Ice Stream bed and those on the Dubawnt Lake Ice Stream bed, Canada (Stokes and Clark, 2003). Figure merged from King *et al.* (2009). [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]

extended along flow  $>10$  km, often with a slightly sinuous nature. Although these features broadly resembled those identified on Rutford Ice Stream (King *et al.*, 2009), Jezek *et al.* (2011) noted that the bed topography and its roughness and structure have the appearance of a bedrock surface with little to no unconsolidated sediment. Calculations of basal drag also indicated a much higher basal shear stress (around 120 kPa) under this ice stream than, for example, those typically found under the Siple Coast ice streams (see section 'West Antarctic ice streams and "Glaciology's grand unsolved problem"'). Thus, the features were interpreted as 'bedrock mega-grooves' that may, in part, be influenced by the underlying bedrock structures, similar to those observed in hard-bed palaeo-ice stream landform assemblages (Smith, 1948; Bradwell, 2005; Bradwell *et al.*, 2008; Eyles, 2012; Krabbendam *et al.*, 2016). If correct, this implies that the features are largely erosional.

More recently, Schroeder *et al.* (2014) used data from radar scattering to image the bed of Thwaites Glacier, West Antarctica. In the upstream region of the ice stream, they were able to detect a corrugated bed with amplitudes ( $\sim 20$  m) and wavelengths ( $\sim 500$  m) that were similar in scale to MSGs found under Rutford Ice Stream (King *et al.*, 2009) and on palaeo-ice stream beds (Livingstone *et al.*, 2012; Spagnolo *et al.*, 2014). However, further downstream, they found topographies that were consistent with a much rougher bed, characterised by outcropping bedrock. This interpretation was also consistent with previous observations of the subglacial water system (Schroeder *et al.*, 2013) and modelling of the

basal shear stresses (Joughin *et al.*, 2009), which found relatively high basal shear stress and channelized subglacial water in the lower trunk region, but with relatively lower basal shear stresses and a distributed water system in the upper trunk and tributaries of the ice stream (see the following section). Elsewhere, on Thwaites Glacier, airborne ice-penetrating radar data aimed at ascertaining the controls on the location of the eastern margin of the ice stream may have inadvertently detected an ice stream shear margin moraine (see Figure 4(a) in MacGregor *et al.*, 2013), although this was not discussed in that study.

### Direct observations of subglacial hydrology and till deltas

The geomorphology associated with subglacial meltwater storage and drainage was also beginning to be uncovered using new remote sensing and geophysical processing techniques. The presence of a distributed system of shallow 'canals' cut into the underlying soft till had been predicted by theory (Alley, 1989a; Ng, 2000; Walder and Fowler, 1994), but evidence of their existence beneath ice streams was very limited on both modern and palaeo-ice stream beds. Carsey *et al.* (2002) had previously drilled into a 1.4 m deep water-filled gap close to the shear margin of Ice Stream C, and Atre and Bentley (1994) interpreted strong reflections  $>50$  km upstream of the

grounding line under Ice Stream B to be ponded water. The pioneering borehole drilling by Engelhardt *et al.* (1990) was also inferred to have punctured an active basal drainage system, but direct evidence of the configuration of subglacial channels or canals remained elusive.

One of the first studies to document the geomorphology associated with meltwater activity under an active ice stream was by King *et al.* (2004), who deployed seismic techniques on Rutford Ice Stream to document the presence of a water-filled canal in deforming till, measuring at least 1 km by 0.2 km and with depths <1 m. King *et al.* (2004) noted that the estimated water layer thickness of their canal was between 0.4 and 0.6 m and that its dimensions were long (> 1 km) and thin (<200 m) and not topographically constrained. Thus, they inferred that they had imaged part of a drainage system, rather than an isolated pond. Later work by Murray *et al.* (2008) used radar and seismic data from the same ice stream to infer several locations where water was interpreted to exist as shallow canals in locations where till was assumed to be deforming. These features were 50 m wide and comprised of water <0.2 m deep. They noted that the features were considerably wider than the 3–5 m predicted by theory (Ng, 2000) or the 1 m inferred by Engelhardt and Kamb (1998), but were narrower than the canal interpreted by King *et al.* (2004).

More recent work by Schroeder *et al.* (2013) on Thwaites Glacier, West Antarctica, also reported substantial volumes of water ponding in distributed canals and increasing in area where ice flow approaches a major bedrock ridge. They used the angular distribution of energy in radar bed echoes to characterise the configuration of subglacial water systems and found that a system of broad distributed channels transitioned into a network of fewer concentrated channels on and over the ridge. The transition between these systems occurred with increasing ice surface slope and basal water flux, which would be predicted by theory (Walder and Fowler, 1994). Schroeder *et al.* (2013) also noted that the transition to a channelized system coincided with an increase in basal shear stress (Joughin *et al.*, 1999), which would also match theoretical predictions.

Elsewhere, the detection of subglacial lakes beneath modern ice masses was a rapidly evolving field (Siegert *et al.*, 2005) and several studies had inferred water bodies (small lakes/ponds) beneath ice streams (Gray *et al.*, 2005; Fricker *et al.*, 2007; Peters *et al.*, 2007; Fricker and Scambos, 2009; Smith *et al.*, 2009). Gray *et al.* (2005), for example, detected ice surface uplift and subsidence on both Kamb Ice Stream (formerly Ice Stream C) and Bindschadler Ice Stream (formerly Ice Stream D), which they attributed to transient movements of subglacial water. The inferred volumetric transfers of water over a 24 day period ( $\sim 20^7$  and  $\sim 10^7$  m<sup>3</sup> on Kamb and Bindschadler, respectively) constituted two major outburst floods of considerable magnitude. Peters *et al.* (2007) also inferred a major meltwater body beneath a tributary of Bindschadler Ice Stream that was trapped by a local reversal in the ice surface slope where the ice stream flowed over rough bed topography. The feature was 5–10 m thick (deep) and at least 1 km long, and existed in a region of the bed with alternating regions of soft and stiff till.

Around the same time, Bindschadler and Choi (2007) suggested that ice stream onset zones would likely be characterised by the presence of subglacial lakes due to the interdependence between rapid ice flow, surface topography and the spatial distribution of subglacial water. Using theories of how subglacial topography is transmitted to the ice surface topography, and how this affects subglacial hydraulic potential and water storage, they posited that the transition from tributary to full ice stream flow closely agrees with where subglacial water may first be stored. It is clear, however, that shallow

(5–10 m) lakes can exist closer to the grounding line, such as Subglacial Lake Whillans (Horgan *et al.*, 2012). A further key study by Fricker *et al.* (2007) documented the presence of several relatively large ( $\sim 50$ – $330$  km<sup>2</sup>) subglacial lakes under Mercer and Whillans Ice Streams (formerly Ice Stream's A and B, respectively) that were associated with ice surface movements of the order of metres over several months, thought to be related to lake volume changes. Later work by Smith *et al.* (2009) also drew attention to the observation that some subglacial lakes in Antarctica appear to be clustered beneath ice streams and other work noted their correspondence to subglacial topography, such as ridges, or areas where there are abrupt contrasts in basal conditions, such as sticky spots (Fricker *et al.*, 2007, 2010; Fricker and Scambos, 2009; Sergienko and Hulbe, 2011). Indeed, Fricker *et al.* (2010) pointed out that the juxtaposition of subglacial lakes and sticky spots is likely due to the fact that the high-friction sticky spots help generate additional meltwater and also cause strong gradients in ice thickness, which feeds back to influence the pattern of hydraulic potential at the bed. This is supported by modelling undertaken by Sergienko and Hulbe (2011), who demonstrated that ice flow over a sticky spot causes favourable conditions for the development of subglacial lakes.

Thus, it was becoming increasingly clear that ponded meltwater, and channels and canals, were perhaps more abundant on ice stream beds than previously recognised (Sergienko and Hulbe, 2011), and there was clear evidence that its water could drain rapidly (Gray *et al.*, 2005; Wingham *et al.*, 2006; Stearns *et al.*, 2008). However, the configuration of the hydrological system during such drainage events was largely unknown. If these drainage events were widely distributed across the bed (i.e. as a thin film or in shallow canals), it could considerably enhance basal lubrication, but flow through a single or small number of larger channels would have much less influence. To assess subglacial water flow to the grounding line of the Siple Coast ice streams, Carter and Fricker (2012) combined lake volume estimates derived from remote sensing with a model for subglacial water transport. They noted that subglacial meltwater outflow tends to concentrate in distinct locations (see also Le Brocq *et al.*, 2013) and that while the mean outflow at the grounding line is around  $60 \text{ m}^3 \text{ s}^{-1}$ , it can increase to  $300 \text{ m}^3 \text{ s}^{-1}$  during synchronised flood events. Carter *et al.* (2017) have also modelled the drainage of lakes in which channels are mechanically eroded into the underlying deformable subglacial sediment. They noted that conventional models based on 'R-channels' cut into the overlying ice are unable to reproduce the timing and magnitude of lake drainage events under ice streams. Their modelling showed how water pressures change during a flood event, such that the initial drainage is associated with a high pressure distributed system, but that this system subsequently collapses to a channelized system. Their modelling was able to replicate both the inferred magnitudes and recurrence intervals of lake-volume changes derived from laser altimeter data on Mercer and Whillans Ice Stream. Release of large volumes of meltwater could also generate significant subglacial erosion and transport, but the extent to which these events do 'geomorphological work' under ice streams is largely unknown (see section 'How is sediment eroded and transported beneath ice streams?').

In relation to sediment transport, geophysical investigations close to the grounding line of active ice streams were now also able to confirm the presence of till deltas that had been hypothesised in the early work by Alley *et al.* (1987a, 1989a) and had subsequently been detected on the beds of numerous marine-based palaeo-ice streams. Anandakrishnan *et al.* (2007) conducted radar surveys in the grounding zone of Whillans Ice Stream, to image a wedge of subglacial sediment with a

maximum thickness of 31 m and extending for >12 km, with the imaged volume on the order of  $10^5 \text{ m}^3$  per unit width (they acknowledged that they were unable to detect the entire feature). Given that the grounding line has likely been in this location for 1000 years (Conway *et al.*, 1999), they calculated a sediment flux of around  $10^2 \text{ m}^3 \text{ m}^{-1} \text{ a}^{-1}$ , which is at the lower end originally predicted by Alley *et al.* (1989a) for the same ice stream ( $10^2$ – $10^3$ ). This rate also implies long-term erosion rates of just over  $0.1 \text{ mm a}^{-1}$  over Whillans Ice Stream and its catchment, although Anandakrishnan *et al.* (2007) noted that erosion is likely to be concentrated in discrete areas where basal melting occurs over poorly consolidated sedimentary substrates. Elsewhere, Doake *et al.* (2001) used a similar approach to Alley *et al.* (1989a) to estimate a higher till flux of  $4 \times 10^3 \text{ m}^3 \text{ m}^{-1} \text{ a}^{-1}$  at the grounding line of Rutford Ice Stream. As in most previous work, the mechanism of sediment transport was assumed to be via mobile layer of deforming sediment tens of centimetres or more, rather than via meltwater and/or melt-out of basal debris. Anandakrishnan *et al.* (2007) drew attention to the similarity between the wedge and grounding zone wedges identified on the sea-floor beyond the limits of the Ross Ice Shelf (Domack *et al.*, 1999; Shipp *et al.*, 1999; Mosola and Anderson, 2006; see section 'Offshore records of palaeo-ice streams based on marine geophysics') and, in a companion paper, Alley *et al.* (2007) demonstrated how such a wedge serves to locally reduce water depths and thereby helps stabilise the ice stream's terminus. In locally stabilising the grounding line, the development of a wedge also allows for focused deposition in that location, rather than the spreading of deposits more uniformly over a broader area (Anandakrishnan *et al.*, 2007).


### Sticky spots and 'traction ribs'

Following the early work that discovered sticky spots beneath active ice streams (see section 'In search of ice stream sticky spots'), new geophysical techniques coupled with inversions of basal drag based on control methods began to uncover spatially organised patterns of high basal drag approximately perpendicular to ice stream flow (MacAyeal, 1992; MacAyeal *et al.*, 1995; Price *et al.*, 2002; Joughin *et al.*, 2004). MacAyeal *et al.* (1995) had first noted bands of higher basal shear stress several tens of kilometres in length and several kilometres in width on Ice Stream E, which they referred to as 'sticky strings'. Price *et al.* (2002) also detected transverse bands of alternating low and high basal shear stress along a tributary of Bindschadler Ice Stream (formerly Ice Stream D), with the bands of high basal shear stress on the order of 10 times the mean values elsewhere. They attributed these bands to the possibility of bedrock ridges striking obliquely across the ice stream and also pointed out that areas of the low shear stress probably contained ponded water in between (see previous section). Joughin *et al.* (2004) subsequently utilised higher resolution velocity datasets to improve the calculations of basal shear stress beneath all of the Siple Coast ice streams. Their results were consistent with previous studies, which indicated a weak bed beneath ice streams, interrupted by localised sticky spots, and with clear patterns of transverse banding apparent, particularly in the tributaries of the ice streams. The cause of this 'banding' in basal shear stress was unknown, but Joughin *et al.* (2004) speculated that they might result from ice flowing across the topographic and tectonic fabric of the underlying terrain, with ice motion accomplished through a combination of soft-bed deformation over sedimentary basins and basal sliding over bedrock ridges. More recently, Peters *et al.* (2007) also noted three broad bands along a 12 km profile in the surface

slope of Bindschadler Ice Stream that were linked to alternating areas of high and low basal shear stress and with water ponding in the areas of low basal shear stress.

More recently, two papers have drawn attention to the apparent ubiquity of these regular patterns of basal shear stress on a number of Antarctic and Greenlandic ice streams, and including non-ice stream areas (Sergienko and Hindmarsh, 2013; Sergienko *et al.*, 2014). These studies utilised higher resolution data on ice velocity, elevation and thickness to calculate basal shear stresses using standard inverse techniques from previous work (MacAyeal, 1992). The increased resolution of these inversions, compared with previous efforts, revealed the presence of regular 'rib-like' patterns of very high basal shear stress (typically ~200–300 kPa) embedded within much larger areas of near-zero basal shear stress (Figure 16). They were found to be widespread throughout areas of slow and fast flow, but they were most pronounced in arcuate patterns within the onset zone of ice streams (Sergienko *et al.*, 2014), as previously noted by Joughin *et al.* (2004). These patterns had a clear surface expression and were seen in the calculations of the driving stress, which are independent of the inversion technique, spatial resolution and its regularisation method. These enigmatic patterns were termed 'traction ribs' (Sergienko and Hindmarsh, 2013) and varied in size from several kilometres to tens of kilometres in length and a few kilometres wide, with the long axes aligned approximately transverse to ice flow, but often deviating by ~30–60° from ice flow direction (Sergienko and Hindmarsh, 2013). It is not clear what causes these regular patterns in basal shear stress, but Sergienko and Hindmarsh (2013) noted the correspondence between the traction ribs and areas of high hydraulic gradient, and suggested that subglacial water may play a role in rib formation. They suggested that the ribs are likely to be regions of variable effective pressure that cause localised strengthening along the base.

In terms of the geomorphology, the resolution of the geophysical data was unable to reveal whether 'traction ribs' had a topographic expression at the bed and whether they were related to an underlying geological control, such as bedrock bumps or ridges (as proposed by Joughin and others, 2004). However, Sergienko and Hindmarsh (2013) noted that their pattern (Figure 16), if not their dimensions, resembled subglacial bedforms observed on palaeo-ice sheet beds, such as the 'mega-ribs' reported by Greenwood and Kleman (2010), and the far more ubiquitous 'ribbed moraines' (Hättestrand, 1997; Hättestrand and Kleman, 1999; Dunlop and Clark, 2006). Despite their similarity in pattern, it was noted that the traction ribs were intermediate in scale between ribbed moraines and mega-ribs and that ribbed landforms at this scale had not been imaged on palaeo-ice streams. Recently, however, Stokes *et al.* (2016b) reported ribbed bedforms from palaeo-ice stream beds in western Canada that resembled both the pattern and scale of the traction ribs reported by Sergienko and Hindmarsh (2013) and Sergienko *et al.* (2014) (see Figure 16(e), (f)). Using Digital Elevation Models, they mapped >1000 rib-like features on four previously-identified palaeo-ice streams from the Interior Plains of Western Canada (Evans *et al.*, 1999, 2008, 2014; Ross *et al.*, 2009; Ó Cofaigh *et al.*, 2010a). Measurements of their length, width, spacing and amplitude showed that they were near-identical to traction ribs, which is consistent with the notion that similar bedforms exist beneath active ice streams and which help explain the transverse banding in basal shear stress. The formation of the ribbed bedforms remains conjectural, but the observations from palaeo-ice streams (Stokes *et al.*, 2016b), coupled with those from modern ice masses and numerical modelling, suggest they might be related to wave-like instabilities occurring in the coupled flow of ice and till and modulated by subglacial meltwater drainage (Dunlop *et al.*, 2008;



**Figure 16.** (a) Inverted basal shear stress for Thwaites Glacier, West Antarctica (in kPa). (b) Observed ice surface speed with black patches ('traction ribs') showing locations where basal shear stress (in a) is  $>100$  kPa. (c) Bed elevation and traction ribs (in black). (d) Gradient of the hydraulic potential (in pascals per meter), with traction ribs (in black). (e) Example of a cluster of low-amplitude rib-like ridges aligned transverse to ice flow on a digital elevation model of the Buffalo Ice Stream corridor in Saskatchewan (see Ross *et al.*, 2009 and Stokes *et al.*, 2016b). (f) Black patches show the location of the central area of rib-like features in (e). (a-d) taken from Sergienko and Hindmarsh (2013; reproduced by permission from The American Association for the Advancement of Science) and (e-f) taken from Stokes *et al.* (2016). [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]

Chapwanya *et al.*, 2011; Sergienko and Hindmarsh, 2013, Fowler and Chapwanya, 2014). Once initiated, high basal shear stresses over the upstanding ridges might also induce glaciotectionism of subglacial sediments, as has been reported for some of the transverse ridges in western Canada (Evans *et al.*, 1999, 2008, 2014).

## Summary

Until the turn of the century, the limited resolution of geophysical techniques had prevented the detailed imaging of the geomorphology of active ice stream beds and much of our understanding of their geomorphology was gleaned from investigations of palaeo-ice stream beds (section 'Identification and Characterisation of the Beds of Palaeo Ice Streams'). However, advances in the resolution and processing of remote sensing and surface and airborne geophysical techniques led to some important discoveries of subglacial bedforms that had previously evaded detection beneath modern ice masses. These studies were able to confirm that many of the landforms first identified on palaeo-ice stream beds, such as MSGs, could be directly linked to ice stream flow (King *et al.*, 2009; Smith

and Murray, 2009). They also confirmed the link between ice velocity and bedform elongation (cf. King *et al.*, 2007), which was initially proposed based only on observations from palaeo-ice stream beds (Clark, 1993). They reinforced the notion that basal conditions are spatially variable, especially in the onset zone of ice streams, and that their motion over a mosaic of sticky and slippery spots is likely accommodated by both till deformation and basal sliding (Smith, 1997a). Higher resolution datasets have also enabled more detailed inversions of basal drag, which revealed a hitherto unrecognised regularity in bed stickiness ('traction ribs') on many ice stream beds (Joughin *et al.*, 2004, Sergienko and Hindmarsh, 2013; Sergienko *et al.*, 2014), which may have a geomorphological expression as ribbed bedforms observed in the palaeo-record (Stokes *et al.*, 2016b). These sticky spots and traction ribs have been shown to be intimately linked to subglacial hydrology (Peters *et al.*, 2007), and the last decade or so has seen an increase in the detection of subglacial lakes (Gray *et al.*, 2005; Peters *et al.*, 2007; Smith *et al.*, 2009) and meltwater drainage systems under active ice streams (King *et al.*, 2004; Schroeder *et al.*, 2013).

Repeat surveying of active ice stream beds (Smith *et al.*, 2007; Smith and Murray, 2009) has also revealed an active

sedimentary system where material is being eroded, transported, and deposited over very short, likely sub-annual time-scales, which is impossible to glean from palaeo-ice stream geomorphology. Moreover, these studies have also allowed estimates of the rates of subglacial erosion. Long-term erosion rates averaged over ice stream catchments have been estimated to be  $0.1\text{--}0.5\text{ mm a}^{-1}$  (Alley *et al.*, 1986; Anadakrishnan *et al.*, 2007), but locally may be as high as  $1\text{ m a}^{-1}$  in some locations (Smith *et al.*, 2007, 2012). Once mobilised, this sediment is thought to be transported downstream, with sediment fluxes at the grounding line typically of the order of  $10^2\text{--}10^3\text{ m}^3\text{ m}^{-1}\text{ a}^{-1}$  (Alley *et al.*, 1989a; Anadakrishnan *et al.*, 2007), but sometimes higher (Doake *et al.*, 2001).

## Future Challenges and Opportunities: Moving from Form to Process in Ice Stream Geomorphology

It is clear that our understanding of ice stream geomorphology has grown rapidly since the early pioneering investigations in West Antarctica in the 1980s (section 'Early Exploration of the Beds of Active Ice Streams'). Initially, our knowledge of the morphology of ice stream beds was largely gleaned from geophysical investigations of modern ice streams in Antarctica. This was soon augmented by observations from palaeo-ice sheets beds and the recognition that ice streams created a distinctive geomorphology that is very different from slower ice sheet flow (Dyke and Morris, 1988). This led to a rapid convergence of ideas and collaboration between glaciology and glacial geomorphology, and went some way to addressing Boulton's (1986) earlier plea for greater interdisciplinarity between these sub-disciplines. Moreover, the last decade or so has seen even greater collaboration between these two sub-disciplines (see also Bingham *et al.*, 2010) and this is largely due to recent technological and methodological advances that have enabled the geomorphology of active ice stream beds to be imaged with unprecedented resolution (Bingham *et al.*, 2010), revealing subglacial bedforms whose interpretation has been considerably aided by their palaeo-counterparts (King *et al.*, 2009; Smith and Murray, 2009). As such, we now have a very detailed knowledge of the morphology of ice stream beds and an emerging understanding of how this relates to the distribution of sediments, water and basal shear stresses. However, our understanding of how the morphology (form) relates to processes is much more limited, and I argue that this represents the next major challenge and will require even closer cooperation between glaciologists, glacial geomorphologists, sedimentologists, together with numerical modelling. For example, the precise mechanisms through which sediment is eroded, transported and deposited under ice streams are largely unknown and remain subject to debate (e.g. deep deforming layer versus shallow deforming layer, channelized meltwater versus canals and water films). These processes are intimately linked to both the growth and decay of subglacial bedforms, but our understanding of bedform creation and evolution is also poorly understood. The way in which ice stream geomorphology influences or even controls ice stream flow over a range of time-scales is also poorly understood, and yet these are a crucial boundary condition of numerical models that aim to predict future ice stream dynamics and their potential contribution to sea level. These issues are now discussed as a series of key questions, together with future opportunities that might enable them to be answered.

## How is sediment eroded and transported beneath ice streams?

The convergence of knowledge from studies of both palaeo- and active ice stream geomorphology indicates that ice streams are responsible for considerable sediment fluxes to their grounding lines, see Table I. As noted by Bougamont and Tulaczyk (2003), this is a key feature of ice streams that must be reproduced by models that attempt to simulate their flow. Moreover, and as noted in section 'Direct observations of subglacial hydrology and till deltas', these high fluxes are important because sediment deposition at ice stream grounding lines can stabilise retreat by serving to locally reduce water depths (Alley *et al.*, 2007). The mechanisms through which sediment is eroded and transported by ice streams is also intimately linked to their basal shear stress, which represents a key boundary condition for numerical modelling of ice stream/ice sheet dynamics (Ritz *et al.*, 2015).

The values in Table I are mostly derived from bracketing the timing of the deposition of large Trough Mouth Fans (Nygård *et al.*, 2007) and grounding zone wedges (Livingstone *et al.*, 2016a), or from modelling sediment transport based on assumed ice and sediment properties (Jenson *et al.*, 1995; Bougamont and Tulaczyk, 2003). Typical sediment fluxes are of the order of  $100\text{ to }1000\text{ m}^3\text{ m}^{-1}\text{ a}^{-1}$  (i.e. per meter width of ice stream terminus), but more extreme values have been estimated from some very large palaeo-ice streams in the Laurentide and Eurasian Ice Sheets. Given these typical values, a key question is: how is the sediment eroded and transported?

Much of the early work invoked viscous deformation of a metres-thick layer of saturated sediment (Alley *et al.*, 1986; Blankenship *et al.*, 1986), which can readily explain the high sediment fluxes and appears to be consistent with observations of both palaeo- (Jenson *et al.*, 1995; Alley, 1991) and modern ice stream sediments and geomorphology (Alley *et al.*, 1987a; Hindmarsh, 1998a, b; Anderson *et al.*, 2002; Dowdeswell *et al.*, 2004; Ó Cofaigh *et al.*, 2005, 2007; Mosola and Anderson, 2006; King *et al.*, 2009; Smith and Murray, 2009). This traditional model has been viewed as the simplest explanation, but it has been repeatedly challenged (and to a large extent been replaced) by evidence that sub-ice stream tills exhibit a near-plastic rheology (Kamb, 1991; Tulaczyk *et al.*, 2000a; Tulaczyk, 2006; Iverson, 2010) and that any deformation should be restricted to a thin zone (of the order of centimetres rather than metres) at the top the till layer and/or accompanied by basal sliding across the surface of the till (Engelhardt *et al.*, 1990; Engelhardt and Kamb, 1998). This is important because if sub-ice stream tills behave as Coulomb-plastic substrates with deformation restricted to narrow shear planes, it potentially limits the thickness of the mobile sediment layer to just a few centimetres and precludes large sediment fluxes via a 'viscously-deforming bed' mechanism (Tulaczyk *et al.*, 2001).

In order to reconcile a Coulomb-plastic rheology with relatively high sediment fluxes (Table I), it is necessary to invoke other mechanisms of sediment transport. One possibility is the ploughing of ice keels through the sediment (Tulaczyk *et al.*, 2001). This mechanism could clearly increase the sediment flux, but generates values (i.e.  $<100\text{ m}^3\text{ m}^{-1}\text{ a}^{-1}$ ) that are at the lower end of those predicted by a viscously-deforming bed model (Tulaczyk *et al.*, 2001). Thus, more recent work has sought to identify other possible sediment transport mechanisms beneath ice streams. Having observed metres-thick layers of frozen sediment on borehole camera imagery from Kamb Ice Stream (formerly Ice Stream C), Christoffersen *et al.* (2010) suggested that basal freeze-on is an important mechanism of sediment entrainment when an

**Table I.** Examples of published estimates of ice stream sediment fluxes

Ice stream (ice sheet)	Sediment flux per metre width of ice stream terminus ( $\text{m}^3 \text{m}^{-1} \text{a}^{-1}$ )	Principal mechanism	Reference
<b>Modern ice streams:</b>			
Ice Stream B (WAIS)	<0.1	<b>Meltwater</b> drainage system	Alley <i>et al.</i> (1989a)
Ice Stream B (WAIS)	< 88	<b>Ploughing</b>	Tulaczyk <i>et al.</i> (2001)
Ice Stream B (WAIS)	~100	<b>Deforming layer</b>	Anandakrishnan <i>et al.</i> (2007)
Ice Stream B (WAIS)	150	<b>Deforming layer</b>	Kamb (2001) and Engelhardt and Kamb (1998), cited in Anandakrishnan <i>et al.</i> (2007)
Ice Stream C (WAIS)	~230	<b>Ploughing</b>	Bougamont and Tulaczyk (2003)
Ice Stream C (WAIS)	525–875	<b>Freeze-on and melt out</b>	Christoffersen <i>et al.</i> (2010)
Ice Stream B (WAIS)	100–1000	<b>Deforming layer</b>	Alley <i>et al.</i> (1989a)
Rutford Ice Stream (WAIS)	4000	<b>Deforming layer</b>	Doake <i>et al.</i> (2001)
<b>Palaeo-ice streams:</b>			
Lake Michigan Lobe (LIS)	~100	<b>Deforming layer</b>	Jenson <i>et al.</i> (1995)
Marguerite Bay palaeo-ice stream (WAIS)	~100–800	<b>Deforming layer</b>	Dowdeswell <i>et al.</i> (2004)
Isfjorden (EIS)	560–980	<b>Deforming layer</b>	Hooke and Elverhøi (1996)
Marguerite Bay palaeo-ice stream (WAIS)	>1000	<b>Deforming layer</b> , plus meltwater and/or freeze-on and melt out	Livingstone <i>et al.</i> (2016a)
Norwegian Channel Ice Stream (EIS)	8000	<b>Deforming layer</b>	Nygård <i>et al.</i> (2007)
Hudson Strait Ice Stream (LIS)	800–37 300	Sediment entrained in <b>basal ice layer</b> (related to Heinrich events)	Dowdeswell <i>et al.</i> (1995)
M'Clintock Channel Ice Stream (LIS)	73 000	<b>Deforming layer</b>	Clark and Stokes (2001)

ice stream shuts-down, and that this sediment can be transported and subsequently released via basal melting when it reactivates. Based on various activation-reactivation scenarios, their estimated sediment fluxes via this mechanism (see Table I) were more comparable with previous estimates that assumed a viscous-bed model (Alley *et al.*, 1989a). Numerical modelling by Bougamont and Tulaczyk (2003) was also able to generate relatively high sediment fluxes (Table I) based on the combined effects of ploughing and the transport of frozen till via plug-flow. Thus, englacial debris transport has potential to be an important component of the relatively high sediment fluxes generated by ice streams (Christoffersen *et al.*, 2010).

High sediment fluxes beneath ice streams might also imply high rates of erosion. Typical steady-state estimates from ice stream catchments averaged over centennial time-scales are generally of the order of  $1 \text{ mm a}^{-1}$ , but there is one example of 1 m of sediment being removed over a just a few years under Rutford Ice Stream (see Table II). Precisely how sediment is eroded is another major area of uncertainty. It has been noted that sub-till erosion by a viscous deforming bed might be too low to sustain high sediment fluxes (Cuffey and Alley, 1996). Indeed, in the absence of other mechanism of erosion to maintain a layer of lubricating sediment, this might make some ice streams susceptible to sediment exhaustion and a transition to

**Table II.** Examples of published estimates of ice stream erosion rates

Ice stream (ice sheet)	Erosion rate ( $\text{mm a}^{-1}$ )	Primary mechanism and context	Reference
Ice Stream C (WAIS)	~0.04–0.1	<b>Basal freeze-on:</b> estimate of the accretion of sediment into a basal ice layer during cycles of basal freeze-on	Christoffersen <i>et al.</i> , (2010)
Ice Stream B (WAIS)	0.1	<b>Deforming layer:</b> long term catchment average	Anandakrishnan <i>et al.</i> (2007)
Ice Stream B (WAIS)	0.1–0.4	<b>Deforming layer:</b> Steady state erosion over catchment	Alley <i>et al.</i> (1987b)
Ice Stream B (WAIS)	~0.5	<b>Deforming layer:</b> Steady state erosion over catchment	Alley <i>et al.</i> (1989a)
Ice Stream C (WAIS)	~0.2–0.6	<b>Basal freeze-on and ploughing:</b> numerical modelling of tributaries and trunk	Bougamont and Tulaczyk. (2003)
Ice Stream B (WAIS)	<0.73	<b>Ploughing:</b> average rate for a 120 km flow-line upstream of UpB Camp	Tulaczyk <i>et al.</i> (2001)
Leverett Glacier (Greenland)	4.8	<b>Meltwater erosion:</b> average rate of erosion from the catchment	Cowton <i>et al.</i> (2012)
Pine Island Glacier (WAIS)	600	<b>Deforming layer:</b> average rate of erosion estimated from ice surface/thickness measurements at one location, averaged over 49 years	Smith <i>et al.</i> (2012)
Rutford Ice Stream (WAIS)	1000	<b>Deforming layer:</b> localised erosion across 500 m wide areas associated with drumlin formation	Smith <i>et al.</i> (2007)

a higher friction bed. Such a process has been invoked to explain the downstream decrease in bedform density and an increase in exposed bedrock on the M'Clintock palaeo-ice stream in Arctic Canada that led to its eventual shut-down (Clark and Stokes, 2001). A similar transition from a very thick (>10m) to much thinner deforming layer (<10m) has also been identified under Pine Island Glacier, West Antarctica (Smith *et al.*, 2013). Given the high erosion rates observed under this ice stream (Smith *et al.*, 2012), Smith *et al.* (2013) hypothesised that the sediment was being progressively eroded and that, if this process continued, the underlying basement rocks might soon outcrop, potentially increasing basal drag and reducing the ice flow.

Sediment freezing on to the base of the ice has been shown to be a particularly effective mechanism of eroding sediment from the bed and into the basal ice layer (Christoffersen *et al.*, 2010). In addition, it has been proposed that basal ice keels might protrude into the sub-strata and cause erosion to replenish a layer of softer till (Tulaczyk *et al.*, 2001; Clark *et al.*, 2003). It is interesting to note, however, that very few studies have invoked erosion and transport via subglacial meltwater (Table II). Meltwater has been shown to preferentially pond in the onset zone of active ice streams (Bindschadler and Choi, 2007; Peters *et al.*, 2007; Schroeder *et al.*, 2013) and there is a wealth of evidence from palaeo-ice stream beds that channelised meltwater flow has incised into bedrock in these locations (Ó Cofaigh *et al.*, 2012; Graham *et al.*, 2009). However, marine geophysical studies have also drawn attention to the fact that there is no evidence for channelized meltwater flow over the soft till that typically characterises fields of MSGs further downstream (Ó Cofaigh *et al.*, 2002; Dowdeswell *et al.*, 2004; Graham *et al.*, 2009). This has often been interpreted to reflect the fact that subglacial meltwater was transported through the soft till matrix and does little geomorphological work, but Ó Cofaigh (2012) pointed out that this may simply reflect the resolution of geophysical techniques. Indeed, canals and ponded water in soft sediment have been detected beneath several West Antarctica ice streams, including small subglacial lakes (Horgan *et al.*, 2012; see also the section 'Direct observations of subglacial hydrology and till deltas') and recent modelling suggests that lake drainage events under ice streams might take place in sediment-floored canals (Carter *et al.*, 2017).

Moreover, recent work from Greenland has suggested that suspended sediment in meltwater emerging from an outlet glacier implies erosion rates as high as  $4.8 \pm 2.6$  mm a year (Cowton *et al.*, 2012), which are much higher than had hitherto been assumed (Table II). Given the emerging evidence of canals and channels beneath some West Antarctic ice streams (King *et al.*, 2004; Schroeder *et al.*, 2013) and that channelized meltwater flow is likely to reach the grounding line of ice streams (Le Brocq *et al.*, 2009, 2013; Dowdeswell and Fugelli, 2012); it seems plausible that subglacial meltwater erosion and transport could play a role in influencing sedimentary processes beneath ice streams and shaping their geomorphology, but one which has been largely overlooked. Indeed, so little work has addressed how meltwater drainage might impact on the geomorphology of ice stream beds that a recent review of the progress in observing and modelling Antarctic subglacial water systems failed to mention the topic (Fricker *et al.*, 2016). In this regard, observations and data from the beds of palaeo-ice streams are a potentially powerful constraint on the configuration of the basal drainage system during lake drainage events. For example, recent work by Livingstone *et al.* (2016b) has documented evidence of the mechanism and geometry of lake drainage events associated with relict subglacial lake locations on the bed of the Laurentide Ice


Sheet. Consistent with work under modern ice streams (see the section 'Direct observations of subglacial hydrology and till deltas'), they showed that the palaeo-lakes were shallow (<10m) lenses of water perched behind ridges orientated transverse to ice flow and that they periodically drained through canals incised into the substrate. The canals were typically 200–300m wide, 4–10m deep and between 700 and 1700m. They also reported that these canals sometimes transition into eskers (formed in R-channels) that likely represent the depositional imprint of the last high magnitude drainage event and supports the notion that subglacial meltwater is a potentially powerful agent of erosion under ice streams.

In summary, determining how sediment is eroded and transported beneath ice streams remains a key challenge. Several key mechanisms have been proposed and these hypotheses are summarised in Figure 17. The next logical step is to attempt to understand which of the processes in Figure 17 are most important and the extent to which they vary both within and between individual ice streams. These issues have attracted the attention of researchers for several decades, but there is little consensus over the primary mechanisms and this fundamentally limits our ability to model ice stream flow and introduces large uncertainties to predictions of their future behaviour. It is clear that sediment fluxes are high and it is likely that a number of mechanisms contribute to sediment erosion and that it is highly variable in space and time (Figure 17). While a large number of studies have invoked erosion and transport associated with a mobile deforming bed, it is possible that the accretion (via basal freeze-on) and subsequent release of sediment from the basal ice layer is also very important. It is also likely that meltwater plays an important role, but this has received very little attention and is a key area for future work to address. Resolving these issues will require major improvements in the resolution of geophysical techniques that can image the beds of active ice streams. Repeat surveys of the bed across larger areas than has hitherto been possible could also be prioritised. Ideally, it would be fruitful to compare surveys from various positions down ice streams (onset, trunk and grounding zone). In addition, *in situ* monitoring of the processes at ice stream grounding lines might also be possible with advances in underwater observing systems. Such data would provide a powerful test for numerical modelling.

## How are bedforms created under ice streams?

The erosion, transport and deposition of sediments under ice streams (Figure 17; section 'How is sediment eroded and transported beneath ice streams?') is intimately linked to the creation of subglacial bedforms. Their growth and evolution introduces basal boundary conditions (e.g. roughness) that are different from a featureless bed and which are likely to modulate the velocity of ice streams (Schoof, 2002). The development of such bedforms is also likely to be intimately linked to subglacial water routing and pressures. However, a further key challenge is that there remains little consensus as to how such bedforms are initiated, evolve and perhaps also decay under ice streams.

As noted above, we now have a very good understanding of what landforms are produced by ice streams (see examples in Figures 5, 9, 12, 15). The most ubiquitous features on ice stream beds are mega-scale glacial lineations (MSGs) (King *et al.*, 2009; Spagnolo *et al.*, 2014). Although most workers have tended to focus on (and map) the ridges, this landscape is perhaps best described as a corrugated surface, and it may be misleading to only focus only the positive relief features (see also the section 'How can we better quantify subglacial



**Figure 17.** Simplified cartoon (not to scale, but resembling the example in Figure 9) illustrating key mechanisms of sediment erosion and transport that have been hypothesised to take place under ice streams. Our understanding of these processes is poorly constrained, particularly with respect to the mechanisms and time-scales of sediment erosion, transport and deposition, and how these relate to the growth and decay of subglacial landforms shown in (a) and the evolution of basal shear stresses. [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]

roughness beneath ice streams?'). Indeed, assemblages of MSGs appear to resemble the scale and pattern of (albeit much rarer) features seen on the bed of ice streams flowing over a largely bedrock substrate and described as mega-grooves (Bradwell, 2005; Bradwell *et al.*, 2008; Jezek *et al.*, 2011; Krabbendam *et al.*, 2016), although this needs testing quantitatively and there are far fewer studies of hard-bedded landform assemblages. Thus, ice streams flowing of a harder bedrock substrate appear to reorganise their bed in a similar fashion to those flowing over soft-sediments, which may point towards mechanisms that lead to net erosion of the subglacial sediments (see Eyles *et al.*, 2016). It is also clear that ice streams that flow through deep troughs (Rutford Ice Stream: King *et al.*, 2009) create MSGs that are near-identical to those on the beds of pure ice streams (the Dubawnt Lake Ice Stream; Stokes *et al.*, 2003). This suggests that although the geometry and force balance of pure and topographic ice streams might be quite different (Bentley, 1987; Truffer and Echelmeyer, 2003), their basal processes and geomorphology might have more in common than previously suggested. Indeed, the only obvious difference between the geomorphology of different types of ice streams is that MSGs on marine-based ice streams are rarely associated with any major meltwater-related features, such as channels, or eskers (Livingstone *et al.*, 2012). In contrast, eskers are commonly reported in association with terrestrially-terminating ice stream beds, but it is often assumed that they were formed during or after ice stream activity (cf. Stokes *et al.*, 2003).

The burgeoning availability of satellite imagery and digital elevation models from both onshore and offshore has enabled workers to collate datasets of several thousand mapped MSGs

and quantitatively characterise their morphometry (Stokes *et al.*, 2013a; Spagnolo *et al.*, 2014). A recent study by Spagnolo *et al.* (2014) analysed around 4000 MSGs from eight ice stream settings: three offshore (Norway and Antarctica), four onshore (Canada), and one from under Rutford Ice Stream, West Antarctica. This revealed that the typical length of MSGs is lower than previously suggested (mode 1000–2000 m; median 2892 m), and that previous work had perhaps tended to over-emphasise the more extreme values of length and elongation ratio. Indeed, while the elongation of MSGs (mode 6–8; median 12.2) is typically higher than features described as drumlins, Spagnolo *et al.* (2014) found that these values, and those of their width (mode 100–200 m; median 268 m), overlap; which suggests the two landforms are part of a morphological continuum and may share a similar origin (see also Stokes *et al.*, 2013a; Ely *et al.*, 2016). They were also able to show that MSGs can initiate and terminate at various locations on an ice stream bed and their exceptional parallel conformity is accompanied by a fairly regular lateral spacing (mode 200–300 m; median 330 m). This, they suggested, provides support to the idea that MSGs are a spatially self-organised phenomenon (see also Spagnolo *et al.*, 2017). They also noted that the size, shape and spatial arrangement of MSGs are consistent both within, and also between, different ice stream beds, and that this is likely to reflect a common mechanism of formation, which is largely insensitive to local factors.

The wealth of data on MSGL size and shape (Stokes *et al.*, 2013a; Spagnolo *et al.*, 2014; Ely *et al.*, 2016) provides a useful test of hypotheses of MSGL formation, but there is no universally-accepted mechanism of their formation. There are

four main ideas that seek to explain their creation: (i) subglacial deformation of till and attenuation downstream (Clark, 1993); (ii) catastrophic meltwater floods (Shaw *et al.*, 2000, 2008); (iii) 'groove-ploughing' by roughness elements (keels) in the basal ice (Tulaczyk *et al.*, 2001; Clark *et al.*, 2003); and (iv) a rilling instability in the basal hydraulic system (Fowler, 2010). Schoof and Clarke (2008) also developed an idea that subglacial flutes may be formed through a transverse secondary flow in basal that can excavate sediment in a cork-screw like fashion and suggested that it might also be applicable to MSGs. However, the generation of secondary flows in ice is not straightforward and they noted that the formation of much wider MSGs would require around 1000 years, which is inconsistent with rapid evolution of bedforms reported under Rutford Ice Stream (Smith *et al.*, 2007; King *et al.*, 2009). The sediment deformation (i) and meltwater flood (ii) hypotheses are extensions of ideas that have been proposed to explain drumlins and, as such, they appeal to the notion of a subglacial bedform continuum (cf. Stokes *et al.*, 2013a; Ely *et al.*, 2016). In contrast, the groove ploughing (iii) and rilling instability (iv) hypotheses appeal to processes that might act to carve a grooved till surface.

The sediment deformation theory is, arguably, seen as most consistent with observations from both modern and palaeo-ice stream beds (Hart, 1997; Hindmarsh, 1998b; King *et al.*, 2009; Smith and Murray, 2009; Clark, 2010). Having formally recognised and named MSGs, Clark (1993) suggested that the extensive literature on other ice-moulded bedforms provided a useful starting point and that the incremental action of ice flow in streamlining MSGL through subglacial deformation/erosion (cf. Boulton, 1987) seemed to be a likely explanation. He argued that if the development of other ice-moulded bedforms, such as drumlins, could initiate by subglacial deformation around inhomogeneities in till, then similar processes might form MSGL, with the difference in scale resulting from increased basal ice velocities. High strain rates, coupled with a plentiful supply of sediment, might lead to subglacial deformation and attenuation of drumlins into much more elongate MSGs. In support of this hypothesis, MSGs are commonly seen to develop downstream of drumlins (Ó Cofaigh *et al.*, 2002; Graham *et al.*, 2009; Stokes *et al.*, 2013a), and sometimes formed side by side with drumlins (Stokes *et al.*, 2013a). They are also predominantly associated with and composed of a 'deformable' till layer (Dowdeswell *et al.*, 2004; Ó Cofaigh *et al.*, 2005, 2007; King *et al.*, 2009; Smith and Murray, 2009). The preferred spacing of MSGs (Stokes *et al.*, 2013a; Spagnolo *et al.*, 2014) is perhaps more difficult to reconcile with initiation from pre-existing inhomogeneities, which are more likely to be randomly located, but recent modelling of the coupled flow of ice, subglacial water and sediment suggests that a regular pattern of 'emergent' MSGL (cf. Clark, 2010) could arise from an instability in the deforming bed (Fowler and Chapwanya 2014). Encouragingly, such modelling is able to make tentative predictions of bedform dimensions, which appear to resemble observations of MSGs, although these comparisons are not straightforward (see discussion in Fowler and Chapwanya, 2014). A potential challenge to the deforming bed hypothesis, however, is that some MSGs are characterised by 'cores' that consist of crudely stratified glaciofluvial sediments, overlain by till (Ó Cofaigh *et al.*, 2013). However, these observations may simply indicate that any subglacial deforming bed must have eroded down into pre-existing sediments (Boyce and Eyles, 1991; Stokes *et al.*, 2013b), unless these sediments were laid down during MSGL formation. In addition, a comprehensive study of MSGs on the bed of a palaeo-ice stream in Poland has shown that their formation likely involves (at least in that location) a process whereby sediment is continuously accreting via a shallow plastically-deforming till layer associated with an inefficient drainage system (Spagnolo *et al.*, 2016).

The meltwater flood hypothesis (Shaw *et al.*, 2008) is largely based on observations of paleo-ice stream geomorphology in Antarctic cross-shelf troughs and invokes catastrophic discharge of turbulent subglacial meltwater. This hypothesis has been applied to drumlin formation (e.g. Shaw, 1983) and large-scale terrestrial flutings (Shaw *et al.*, 2000) and is based largely on form analogy between MSGs and similar bedforms and patterns created by broad, turbulent flows in water and air. The form analogy is persuasive and Shaw *et al.* (2008) also point to the abundance of meltwater features and tunnel channels in ice stream onset zones (e.g. crescentric and hairpin scours around the stoss end of drumlins and MSGL), and numerous gullies and channels that often characterise the continental slope. However, there is a general absence of meltwater channel features either upstream or downstream of the MSGs on most ice stream beds and there is the long-standing issue as to whether the magnitude of meltwater required to form such floods is plausible (Clarke *et al.*, 2005; Ó Cofaigh *et al.*, 2010b). It is also the case that there have been no observations of significant meltwater floods during the formation of bedforms under Rutford Ice Stream (Smith and Murray, 2009; King *et al.*, 2009), which Shaw and Young (2010) acknowledged would 'oblige us to take a long, hard look at the megaflood hypothesis' (p. 199). As noted, however, meltwater drainage events have been observed under some ice streams (see the section 'Direct observations of subglacial hydrology and till deltas') and a key area for future work is to ascertain the nature and form of these events and the likely impact on sub-ice stream geomorphology (Livingstone *et al.*, 2016b).

The groove-ploughing hypothesis invokes roughness elements in the basal ice (keels) that are able to plough through soft sediments and excavate grooves which leave MSGs as erosional remnants (Clark *et al.*, 2003). An important assumption with the groove-ploughing mechanism is that grooves can plough through sediment for a sufficient distance (several kms) without thermodynamic and mechanical degradation. Clark *et al.* (2003) argued that larger keels (30 m wavelength, 5 m amplitude) are more likely to survive (Thorsteinsson and Raymond, 2000) and that survival distances of 10–100 km are plausible, depending on ice velocity (Thorsteinsson and Raymond, 2000; Tulaczyk *et al.*, 2001; Clark *et al.*, 2003). Clark *et al.* (2003) outlined several predictions of the groove-ploughing hypothesis. An obvious implication is that the transverse roughness of ice stream beds should greatly exceed explicitly measured or quantified roughness, which has been observed in numerous studies (Siegert *et al.*, 2004; Bingham and Siegert, 2007, 2009). A further prediction is that MSGs should occur in areas downstream of regions where basal roughness is produced (e.g. downstream of zones of flow convergence and/or more resistant bedrock), and this has also received much observational support (Ó Cofaigh *et al.*, 2002; Graham *et al.*, 2009; Livingstone *et al.*, 2012). Clark *et al.* (2003) also predicted that the transverse groove spacing should be related to the spatial frequency of roughness that is generated upstream, but MSGs have been observed to initiate within existing 'grooves' (Stokes *et al.*, 2013a; Spagnolo *et al.*, 2014); and the finding that MSGs exhibit a preferred lateral spacing (Stokes *et al.*, 2013a; Spagnolo *et al.*, 2014) is difficult to reconcile with this prediction. In addition, and where it has been explicitly measured (Stokes *et al.*, 2013a; Spagnolo *et al.*, 2014), there is little indication that groove width and depth decrease in the downstream direction, which was also predicted by Clark *et al.* (2003). Thus, while there is some observational support for groove-ploughing and it is intuitively attractive, it does not appear to be the primary mechanism through which MSGs are formed.

The rilling instability theory was put forward by Fowler (2010) and is based on the theory that a uniform water-film

(~2–3 mm deep) flowing between ice and deformable subglacial till would be unstable and rilling would occur, similar to that seen in subaerial settings and which results in a number of narrow streams separated by intervening ridges. Fowler (2010) numerically modelled this process and, for a particular choice of parameters used, he found a preferred spacing (distance to nearest lateral neighbour: 394 m), length scale (52.9 km) and amplitude (12.3 m) which were broadly consistent with observations (Spagnolo *et al.*, 2014). However, these results were strongly dependent on the choice of parameters selected. The appeal of this mechanism is that it can simulate the regularity and spatial organisation of MSGs (Spagnolo *et al.*, 2014; 2017). Moreover, this process has since been incorporated into a more sophisticated treatment that combines both fluvial sediment transport and the coupled flow of ice and sediment to produce a continuum of ribbed moraine, drumlins and MSGs (Fowler and Chapwanya, 2014).

In summary, there is no consensus as to how MSGs are generated beneath ice streams and how the processes summarised in Figure 17 might contribute to their formation. Given the available observations of both the geomorphology and composition of MSGs, it seems likely that their formation primarily involves a mobile layer of deforming till (albeit of uncertain thickness) that is transported/attenuated downstream (cf. King *et al.*, 2009) and which probably involves net erosion of the subglacial sediments over long (> centennial) time-scales. A further appeal of this as a primary mechanism is that it may also explain the broader continuum of landforms found on ice stream beds, including drumlins and ribbed moraines, and the recently inferred 'traction ribs' (section 'Sticky spots and "traction ribs"'). However, given that the pervasive and viscous deformation of several metres of till has largely fallen out of favour in recent years, it is probable that localised erosion from both ice keel ploughing and meltwater might also take place with grooves, in addition to basal freeze-on and subsequent melt-out. These mechanisms might also help to reconcile the high rates of sediment erosion and transport beneath ice streams (see Tables I and II), but their relative roles remains largely unknown. In terms of other bedforms, ice stream shear margin moraines have received minimal attention in the literature (Stokes and Clark, 2002b; Hindmarsh and Stokes, 2008; Batchelor and Dowdeswell, 2016), and their origin is even more enigmatic, despite their potential role in helping to stabilise lateral shear margins and the width of ice streams. They represent a key area for future work to address, in addition to a more quantitative comparison between soft-bedded MSGs and their qualitatively similar bedrock mega-grooves.

### How can we better quantify subglacial roughness beneath ice streams?

Studies of ice stream geomorphology (Stokes *et al.*, 2013a) have perhaps been preoccupied with mapping landforms and quantifying the size and shape of individual features. This is probably because the early work on palaeo-ice streams was simply trying to identify their location and incorporate them into a palaeo-glaciological reconstruction (see the section 'In Search of palaeo-ice streams'). It soon became clear that extracting 'metrics' from these mapped features could lead to new insights with regard to the subglacial landscape and the formation of subglacial bedforms (see previous section), but there are other means of quantifying ice stream geomorphology that have perhaps been under-utilised, particularly with respect to subglacial roughness and how flow-sets of bedforms evolve through time.

Bingham *et al.* (2010) defined bed roughness as the vertical variation of an ice-sheet bed with horizontal distance and

noted that it can be quantified in various ways. Radar-derived ice bed topography has been used most widely to investigate the roughness of ice stream beds and one approach has been to simply characterise roughness based on the standard deviations of individual bed elevation points from an interpolated bed surface (Rippin *et al.*, 2006). More sophisticated techniques involve assessing bed roughness based on the power spectra derived from Fast-Fourier-transforming bed-elevation profiles (Taylor *et al.*, 2004), which has been used to quantify bed roughness across several West Antarctic ice streams (Siegert *et al.*, 2004; Bingham and Siegert, 2007; Bingham and Siegert, 2009). Taken together, these have clearly shown that the bed roughness beneath ice streams is much smoother than beneath non-streaming ice and that roughness also tends to decrease down ice streams.

Bingham and Siegert (2009) usefully outlined a framework for the geomorphological interpretation of bed roughness in Antarctica, suggesting that the smoother beds of ice streams likely result from pre-existing marine sediments, together with warm-based ice flowing at high velocities, which promotes subglacial erosion and deposition. They suggested that it would be worthwhile to apply parallel methods to former ice sheet beds in order to develop a methodological framework that captures the signature of roughness in formerly glacial landscapes, and this would seem an obvious priority for future work. That is, methods could be developed to quantify the roughness of various assemblages of landforms on palaeo-ice stream beds. If it can be shown that different landform assemblages have different roughness signatures, then this opens up huge potential to use measurements of roughness on modern ice streams to infer distinct geomorphologies that may lie at their bed. To that end, Fourier analysis was recently used to analyse MSGs from a number of different ice stream beds, which is one of the first attempts to quantify and compare MSG topographies, but without the need for mapping individual bedforms (Spagnolo *et al.*, 2017). The results from that study concluded that assemblages of MSGs were very similar across different settings, perhaps reflecting pattern evolution via downstream wavelength coarsening, and consistent with the instability theory for subglacial bedforms (Fowler and Chapwanya, 2014).

It would also be useful to quantify how roughness changes along a flow-line from the slow-flowing ice (close to the ice divide), through ice stream onset zones, and into the main trunk, and see how this compares with ice velocities. Is there, for example, a threshold roughness that dictates the location of ice stream onset? Does the roughness on a soft-bedded ice stream differ from a hard-bedded ice stream? Can soft-bedded ice streams transition into hard-bedded ice streams through a process of sediment exhaustion? Another key area of investigation would be to measure and quantify how values of roughness change through time. A fundamental and yet largely unanswered question is: do ice stream beds get rougher or smoother and over what time-scales (or can they do both, depending on underlying geology and setting)? This could be measured on active ice streams over short (annual to decadal time-scales), but it might also be possible to extract measurements of roughness from palaeo-ice stream beds that are well-dated in terms of their duration. The quantification of 'mature' versus 'younger' ice stream geomorphology would provide a useful framework for interpreting the age of ice stream beds in both modern and palaeo-ice sheets. Moreover, if it can be shown that roughness evolves through time in a predictable manner, this would help parameterise the evolution of basal shear stress in numerical ice sheet models that aim to predict future ice sheet dynamics.

Related to the issue of roughness is the recent analysis of the size–frequency distribution of subglacial bedforms in flow-sets. Large datasets of bedform metrics have revealed that drumlins

and MSGs display smooth unimodal frequency distributions with a positive skew (Clark *et al.*, 2009; Stokes *et al.*, 2013a; Spagnolo *et al.*, 2014). When their length, amplitude or spacing are converted to their natural logarithm, the resulting size–frequency distribution is well-defined as log-normal (Fowler *et al.*, 2013; Hillier *et al.*, 2013; Spagnolo *et al.*, 2014). This distribution is extremely common in nature and is thought to emerge from a large number of independent events in which incremental growth (or decay) can occur (Fowler *et al.*, 2013; Spagnolo *et al.*, 2014). Thus, several studies have argued that the observed log-normal distributions of MSGs (Spagnolo *et al.*, 2014) implies an element of randomness in their development, such that growth phases occur randomly and for random duration, likely alongside episodes of decay, e.g. perhaps associated with episodes of bedform erosion and deposition (Fowler *et al.*, 2013; Spagnolo *et al.*, 2014; Hillier *et al.*, 2013, 2016). As noted by Hillier *et al.* (2013), this is consistent with the geophysical studies that have revealed variable bed conditions in both space and time (Smith, 1997a, b; Murray *et al.*, 2008) and subglacial landforms (King *et al.*, 2007; Smith and Murray, 2009) that evolve rapidly on sub-decadal time-scales (Smith *et al.*, 2007; King *et al.*, 2009). Hillier *et al.* (2013) also noted, however, that it is unclear whether this variability arises from the dynamics of ice–sediment–water interactions (e.g. basal stick–slip events) or from interactions between bedforms. Notwithstanding this uncertainty, this stochastic approach contrasts with a deterministic view whereby proto-bedforms of known size and shape always evolve similarly with time to a predictable final morphology. A key challenge, therefore, is to try and incorporate these elements of stochasticity into models of bedform genesis (see next section).


Similar to the analyses of roughness outlined above, Hillier *et al.* (2013) also noted that it would be useful to ascertain the stability or otherwise of bedform populations through time,

i.e. are they in steady state? Using observations of flow-sets beneath modern ice streams, it might also be possible to create statistical models that link the physical processes to observable characteristics of bedform populations under ice streams (see Hillier *et al.*, 2016).

### Can we numerically model ice stream geomorphology?

As noted in the section ‘How are bedforms created under ice streams?’, there are several competing hypotheses that seek to explain ice stream geomorphology, almost exclusively focusing on the formation of MSGs. Unfortunately, only very rarely have these ideas developed to the stage where they can be captured in physically-based numerical models of the ice–bed interface, and this is a further key challenge for future work to address. Of the various hypotheses relating to ice stream geomorphology, only the deforming bed/till instability theory has seen some success in being able to produce a recognisable geomorphology (Hindmarsh, 1998a, b; Schoof, 2007; Chapwanya *et al.*, 2011; Fowler and Chapwanya, 2014). The advantage of these models being able to predict ice stream geomorphology is that they can be tested against large datasets of landform metrics that now exist (Spagnolo *et al.*, 2014), although this has proved difficult, except for ribbed moraine (Dunlop *et al.*, 2008; Chapwanya *et al.*, 2011). The ultimate aim would be for a model to be able to simulate the formation of landform assemblages that resemble those on ice stream beds, much like Dunlop *et al.* (2008) were able to simulate modelled ribbed moraines that matched observations.

Recently, Barchyn *et al.* (2016) took a reduced complexity approach in order to try and simulate recognisable bedforms



**Figure 18.** Samples of output from a reduced complexity model aimed at replicating the initiation and development of subglacial bedforms under ice streams (from Barchyn *et al.*, 2016). The transition between ribbed moraines (a), drumlins (b) and mega-scale glacial lineations (c) is associated with increasing ice speeds and declining sediment thickness. Reproduced by permission of John Wiley and Sons Ltd. [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]

using simple mechanisms based on shallow sediment dynamics that are hypothesised to exist subglacially (Figure 17). Although their focus was on subglacial bedforms in general, rather than specifically on ice stream geomorphology, they found that bedforms readily emerge, see Figure 18, and their interactions mirrored those in aeolian and fluvial geomorphology. Despite its reductionist approach, they were also able to replicate a bedform continuum, whereby transitions between ribbed moraines and elongate flow-parallel bedforms were associated with increasing ice speeds and decreasing sediment thickness. In their model, drumlins transition into MSGLs because the lower ice-to-bed pressure and high ice velocities force bedforms to elongate by extending cavities and low-pressure areas on the lee-side of bedforms. This model was also able to predict till fluxes (of the order of  $10\text{ m}^3\text{ m}^{-1}\text{ a}^{-1}$  for ice flowing at  $150\text{ m a}^{-1}$ ), which they noted was similar but perhaps an underestimate compared with values elsewhere. As in most previous modelling efforts, the aim was to try to replicate the characteristic dimensions of subglacial bedforms (lengths, widths, heights) and sediment properties and processes were necessarily simplified. Clearly, as these modelling experiments evolve, they will probably benefit from more sophisticated parameterisations based on known sediment properties of the ice stream bedforms.

The value of these modelling approaches is that observable characteristics of bedforms (their geomorphology) can then be securely linked to processes and the properties of the overlying ice (thickness, velocity, etc.). Not only would this help inversions of ice sheet dynamics from palaeo-ice stream geomorphology, but it would also allow forward runs to observe the time-scales over which the geomorphology evolves and influences basal shear stresses. Such knowledge is required to reduce the uncertainties in numerical models of future ice sheet/ice stream dynamics, where the parameterisation and evolution of basal shear stresses is a key unknown (Ritz *et al.*, 2015).

In developing the next generation of numerical models of sub-ice stream processes, it is useful to consider what key aspects of ice–bed interface (Figure 17) should be targeted by a successful model. Based on this review of the literature, it seems that the following, at least in part, are salient features of the subglacial environment of ice streams:

- The efficient transport of sediment downstream to the grounding line to account for high sediment fluxes and the building of grounding zone wedges (essential)
- The creation of highly elongate subglacial bedforms (MSGLs) that elongate with ice velocity (essential) and show transitions from drumlins to MSGLs (desirable)
- A mechanism (or mechanisms) to erode and replenish subglacial sediment (desirable)
- A mechanism (or mechanisms) that generates spatial and temporal variability in basal shear stress (desirable)

## Conclusions

Rapidly-flowing ice streams are an important component of ice sheet mass balance and associated impacts on sea level. However, it was only in the 1970s that their importance was fully recognised and the study of ice streams began in earnest. It soon became clear that their flow was governed by processes at the ice–bed interface that was characterised by a unique geomorphology. Elucidating how this geomorphology is created and evolves is critical to understanding and modelling ice stream flow, and our knowledge has grown rapidly over the last three decades, from almost complete ignorance to a detailed knowledge of the morphology of ice stream beds. This

has been brought about through: (i) geophysical investigations of active ice streams, mostly in the West Antarctica Ice Sheet; and through (ii) the investigation of sediments and landforms left behind by ice streams in areas formerly occupied by ice sheets. This paper has reviewed progress in these two main areas from a historical perspective in order to identify key areas of progress in both glaciology and glacial geomorphology and how they have been brought about. Emphasis has been placed on the extent to which these sub-disciplines have converged to help understand ice stream geomorphology, together with the key challenges that remain and how they might be overcome.

From this review, it is clear that we now have a very detailed knowledge of the morphology of ice stream beds and an emerging understanding of how this relates to the distribution of sediments, water and basal shear stresses. However, our understanding of how the morphology (form) relates to processes is much more limited, and I argue that this represents the next major challenge and will require even closer cooperation between glaciologists, glacial geomorphologists, and sedimentologists, together with numerical modellers. Knowledge from both palaeo and modern ice stream studies has allowed for improved estimates of the rates of subglacial erosion and transport (Tables I and II), but the importance of various mechanisms of sediment erosion and transport (Figure 17) is largely unknown, and processes involving subglacial meltwater are potentially important, but have perhaps been overlooked in terms of shaping sub-ice stream geomorphology. Furthermore, sediment erosion, transport and deposition are likely to be intimately linked to the growth and decay of subglacial bedforms, which modulate the velocity of ice streams through their influence on basal roughness and subglacial hydrology. Unfortunately, only very rarely have formational hypotheses (e.g. of MSGLs) been developed to the stage where they can be captured in physically-based numerical models of the ice–bed interface, and this is a further key challenge for future work to address. Such modelling is likely to benefit from the integration of the known properties of subglacial sediments, which are readily accessible on palaeo-ice stream beds but have perhaps been under-utilised.

The way in which ice stream geomorphology influences or even controls ice stream flow over a range of time-scales is also poorly understood, and yet such knowledge is required for numerical modelling that aims to predict future ice stream dynamics and their potential contribution to sea level. Studies of ice stream geomorphology have perhaps been preoccupied with mapping landforms and quantifying the size and shape of individual features (object-based morphometry). Quantification of these ‘metrics’ has led to some important insights with regard to the formation of MSGLs, but other means of quantifying ice stream geomorphology have perhaps been under-utilised, particularly with respect to how topography (or roughness) can be quantified without the need to ‘map’ individual features (general geomorphometry). In addition, and with repeat surveys of ice stream beds now being undertaken (King *et al.*, 2017), it should be possible to examine how fields (or flow-sets) of bedforms evolve through time, which could be explored by statistical analysis of size–frequency distributions. These methods of quantifying sub-ice stream geomorphology also offer a tangible means to help parameterise the evolution of basal shear stress in numerical ice sheet models and there is room for much closer collaboration with those modelling bedforms and basal shear stresses in aeolian and fluvial environments. Given the increasing resolution of digital elevation models from both palaeo and modern ice stream settings, there are several key questions that should be tractable within a few years: is there a threshold roughness that dictates the onset of ice stream flow? Do ice stream beds get rougher or smoother and over what time-scales? To what extent is the

geomorphology on hard and soft-bedded ice streams different, and how might this influence their flow?

In summarising his thoughts on the Cordilleran Ice Sheet Symposium in 1990 and noting the ‘currently fashionable term ‘ice stream’, Mathews (1991: p. 265) stated that ‘with so little known about the conditions and processes operating at the bed of contemporary ice streams, it seems doubtful that the site of an ancient ice stream can be identified solely from a track engraved on the substratum. It will remain a challenge until more is learned about the subglacial topography of the Greenland and Antarctic ice sheets and the erosional processes under both slow and fast-moving ice’. It is pleasing to see that three decades of collaboration between both glaciologists and glacial geomorphologists has overcome this challenge and we now have a detailed knowledge of the glacial geomorphology engraved by ice streams. However, there remains a challenge to learn more about the processes that create ice stream geomorphology and how they impact on ice dynamics. The next three decades are likely to see further growth in the imaging of geomorphology under active ice streams, which should help meet this challenge and provide the necessary constraints for numerical models that are urgently required in the context of ice sheet stability in a warming climate.

**Acknowledgements**—I am indebted to my PhD supervisor Chris Clark who was among the first to recognise the importance of identifying the ‘footprints’ of palaeo-ice streams and who ignited my interest in ice stream geomorphology. Many of the ideas in this paper have also been provoked and nourished by discussions and collaboration with numerous individuals who have always been generous with their time and insights on this subject. In addition to Chris, I would particularly like to thank Johan Kleman, Slawek Tulaczyk, Olav Lian, Richard Hindmarsh, Andrew Fowler, Karin Andreassen, Monica Winsborrow, Dag Ottesen, Ed King, Matteo Spagnolo, Stephen Livingstone and Jeremy Ely, together with my colleagues in Durham: Colm Ó Cofaigh, Dave Evans, Dave Roberts, and Stewart Jamieson. I would also like to thank Chris Orton, Department of Geography at Durham University, for drawing some of the Figures. The Centre for Arctic Gas Hydrate, Environment and Climate (CAGE), at the Department of Geology at UiT, the Arctic University of Norway hosted me as a Visiting Researcher during the completion of this manuscript, for which I am very grateful. I would also like to acknowledge funding from the European Union Initial Training Network (EU ITN) ‘GLANAM’ Project (317217, FP7 2007/2013) and NERC grant NE/J00782X/1. I would also like to thank the Editor – Stuart Lane – for inviting me to write this review, the Associate Editor, and, in particular, Andy Smith, whose comments during the peer-review process were invaluable.

## References

- Alley RB. 1989a. Water-pressure coupling of sliding and bed deformation: I. Water system. *Journal of Glaciology* **35**(119): 108–118.
- Alley RB. 1989b. Water-pressure coupling of sliding and bed deformation: II. Velocity-depth profiles. *Journal of Glaciology* **35**(119): 119–129.
- Alley RB. 1991. Deforming bed origin for southern Laurentide till sheets? *Journal of Glaciology* **37**(125): 67–76.
- Alley RB. 1993. In search of ice stream sticky spots. *Journal of Glaciology* **39**(133): 447–454.
- Alley RB. 2000. Continuity comes first: recent progress in understanding subglacial deformation. In *Deformation of Glacial Materials*, Maltman AJ, Hubbard B, Hambrey MJ (eds), Vol. **176**. The Geological Society of London Special Publication; 171–179.
- Alley RB, Anandakrishnan S, Dupont TK, Parisek BR, Pollard D. 2007. Effect of sedimentation on ice-sheet grounding-line stability. *Science* **315**: 1838–1841.
- Alley RB, Blankenship DD, Bentley CR, Rooney ST. 1986. Deformation of till beneath ice stream B, West Antarctica. *Nature* **322**: 57–59.
- Alley RB, Blankenship DD, Bentley CR, Rooney ST. 1987a. Till beneath Ice Stream B: 3. Till deformation: evidence and implications. *Journal of Geophysical Research* **92**(B9): 8921–8929.
- Alley RB, Blankenship DD, Rooney ST, Bentley CR. 1987b. Till beneath Ice Stream B: 4. A coupled ice-till flow model. *Journal of Geophysical Research* **92**(B9): 8931–8940.
- Alley RB, Blankenship DD, Rooney ST, Bentley CR. 1989a. Sedimentation beneath ice shelves – the view from Ice Stream B. *Marine Geology* **85**: 101–120.
- Alley RB, Blankenship DD, Rooney ST, Bentley CR. 1989b. Water-pressure coupling of sliding and bed deformation: III. Application to Ice Stream B, Antarctica. *Journal of Glaciology* **35**(119): 130–139.
- Alley RB, Clark PU, Huybrechts P, Joughin I. 2005. Ice sheet and sea level changes. *Science* **310**: 456–460.
- Alley RB, Lawson DE, Larson GJ, Evenson EB, Baker GS. 2003. Stabilizing feedbacks in glacier bed erosion. *Nature* **424**: 758–760.
- Allison I. 1979. The mass budget of the Lambert Glacier drainage basin, Antarctica. *Journal of Glaciology* **22**(87): 223–235.
- Anandakrishnan S, Alley RB. 1994. Ice Stream C, Antarctica, sticky spots detected by microearthquake monitoring. *Annals of Glaciology* **20**: 183–186.
- Anandakrishnan S, Alley RB. 1997. Stagnation of Ice Stream C, West Antarctica by water piracy. *Geophysical Research Letters* **24**(3): 265–268.
- Anandakrishnan S, Bentley CR. 1993. Micro-earthquakes beneath ice streams B and C, West Antarctica: observations and implications. *Journal of Glaciology* **39**(133): 455–462.
- Anandakrishnan S, Catania GA, Alley RB, Horgan HJ. 2007. Discovery of till deposition at the grounding line of Whillans Ice Stream. *Science* **315**: 1835–1838.
- Anderson JB, Shipp SS, Lowe AJ, Wellner JS, Mosola AB. 2002. The Antarctic Ice Sheet during the Last Glacial Maximum and its subsequent retreat history: a review. *Quaternary Science Reviews* **21**: 49–70.
- Andrews JT. 1982. On the reconstruction of Pleistocene ice sheets: a review. *Quaternary Science Reviews* **1**(1): 1–30.
- Atre SR, Bentley CR. 1993. Laterally varying basal conditions beneath Ice Streams B and C, West Antarctica. *Journal of Glaciology* **39**(133): 507–514.
- Atre SR, Bentley CR. 1994. Indication of a dilatant bed near Downstream B Camp, Ice Stream B, Antarctica. *Annals of Glaciology* **20**: 177–182.
- Aylsworth JM, Shilts WW. 1989. Glacial Features around the Keewatin Ice Divide: Districts of Mackenzie and Keewatin. *Geological Survey of Canada Paper* **88–24**: 21.
- Bamber JL, Vaughan DG, Joughin I. 2000. Widespread complex flow in the interior of the Antarctic Ice Sheet. *Science* **287**(5456): 1248–1250.
- Barchyn TE, Dowling TPF, Stokes CR, Hugenholtz CG. 2016. Subglacial bedform morphology controlled by ice speed and sediment thickness. *Geophysical Research Letters* **43**. <https://doi.org/10.1002/2016GL069558>.
- Batchelor CL, Dowdeswell JA. 2014. The physiography of High Arctic cross-shelf troughs. *Quaternary Science Reviews* **92**: 68–96.
- Batchelor CL, Dowdeswell JA. 2015. Ice-sheet grounding-zone wedges (GZWs) on high-latitude continental margins. *Marine Geology* **363**: 65–92.
- Batchelor CL, Dowdeswell JA. 2016. Lateral shear-moraines and lateral marginal-moraines of palaeo-ice streams. *Quaternary Science Reviews* **151**: 1–26.
- Bell R. 1895. The Labrador Peninsula. *Scottish Geographical Magazine* **11**: 335–361.
- Benn DI, Evans DJA. 2010. *Glaciers and Glaciation*, 2nd edn. London UK: Hodder Education.
- Bennett MR. 2003. Ice streams as the arteries of an ice sheet: their mechanics, stability and significance. *Earth-Science Reviews* **61**: 309–339.
- Bentley CR. 1976. High electrical-resistivity deep in Antarctic shelf ice or ice stream origin. *Transactions of the American Geophysical Union* **57**(4): 243–243.
- Bentley CR. 1987. Antarctic ice streams: a review. *Journal of Geophysical Research* **92**(B9): 8843–8858.
- Bindschadler RA, Choi H. 2007. Increased water storage at ice-stream onsets: a critical mechanism? *Journal of Glaciology* **53**(181): 163–171.
- Bindschadler RA, Scambos TA. 1991. Satellite-image-derived velocity field of an Antarctic ice stream. *Science* **252**(5003): 242–246.

- Bindschadler RA, Stephenson SN, MacAyeal DR, Shabtaie S. 1987. Ice dynamics at the mouth of Ice Stream B, Antarctica. *Journal of Geophysical Research* **92**(B9): 8885–8894.
- Bingham RG, King EC, Smith AM, Pritchard HD. 2010. Glacial geomorphology: towards a convergence of glaciology and geomorphology. *Progress in Physical Geography* **34**(3): 327–355.
- Bingham RG, Siegert MJ. 2007. Bed roughness characterization of Institute and Möller Ice Streams, West Antarctica: comparison with Siple Coast ice streams. *Geophysical Research Letters* **34**: L21504, <https://doi.org/10.1029/2007GL031483>.
- Bingham RG, Siegert MJ. 2009. Quantifying subglacial bed roughness in Antarctica: implications for ice-sheet dynamics and history. *Quaternary Science Reviews* **28**: 223–239.
- Blake E, Clarke GKC, Gérin MC. 1992. Tools for examining subglacial bed deformation. *Journal of Glaciology* **38**(130): 388–396.
- Blankenship DD, Bentley CR, Rooney ST, Alley RB. 1986. Seismic measurements reveal a saturated porous layer beneath an active Antarctic ice stream. *Nature* **322**: 54–57.
- Blankenship DD, Bentley CR, Rooney ST, Alley RB. 1987. Till beneath Ice Stream B. 1. Properties derived from seismic travel times. *Journal of Geophysical Research* **92**(B9): 8903–8911.
- Bluemle JP, Lord LM, Hunke NT. 1993. Exceptionally long, narrow drumlins formed in subglacial cavities, North Dakota. *Boreas* **22**: 15–24.
- Bougamont M, Tulaczyk S. 2003. Glacial erosion beneath ice streams and ice-stream tributaries: constraints on temporal and spatial distribution of erosion from numerical simulations of a west Antarctic ice stream. *Boreas* **32**: 178–190.
- Bougamont M, Tulaczyk S, Joughin I. 2003a. Response of subglacial sediments to basal freeze-on: 2. Application in numerical modelling of the recent stoppage of Ice Stream C, west Antarctica. *Journal of Geophysical Research* **108**(B4): 20.1–20.16.
- Bougamont M, Tulaczyk S, Joughin I. 2003b. Numerical investigations of the slow-down of Whillans Ice Stream, west Antarctica: is it shutting down like Ice Stream C? *Annals of Glaciology* **37**: 239–246.
- Boulton GS. 1976. The origin of glacially fluted surfaces - observations and theory. *Journal of Glaciology* **17**: 287–309.
- Boulton GS. 1986. A paradigm shift in glaciology? *Nature* **322**: 18.
- Boulton GS. 1987. A theory of drumlin formation by subglacial sediment deformation. In *Drumlin Symposium*, Menzies J, Rose J (eds). A.A. Balkema: Rotterdam; 25–80.
- Boulton GS, Clark CD. 1990. A highly mobile Laurentide Ice Sheet revealed by satellite imagery of glacial lineations. *Nature* **346**(6287): 813–817.
- Boulton GS, Dobbie KE, Zatsepin S. 2001. Sediment deformation beneath glaciers and its coupling to the hydraulic system. *Quaternary International* **86**: 3–28.
- Boulton GS, Hindmarsh RCA. 1987. Sediment deformation beneath glaciers: rheology and geological consequences. *Journal of Geophysical Research* **92**(B9): 9059–9082.
- Boulton GS, Jones AS. 1979. Stability of temperate ice caps and ice sheets resting on beds of deformable sediment. *Journal of Glaciology* **24**(90): 29–43.
- Boyce JJ, Eyles N. 1991. Drumlins carved by deforming till streams below the Laurentide ice sheet. *Geology* **19**: 787–790.
- Bradwell T. 2005. Bedrock megagrooves in Assynt, NW Scotland. *Geomorphology* **65**(3–4): 195–204.
- Bradwell T, Stoker M, Krabbendam M. 2008. Megagrooves and Streamlined Bedrock in NW Scotland: The role of ice streams in landscape evolution. *Geomorphology* **97**(1–2): 135–156.
- Briner JP. 2007. Supporting evidence from the New York drumlin field that elongate subglacial bedforms indicate fast ice flow. *Boreas* **36**(20): 143–147.
- Brookes IA. 2007. First recognition of a Laurentide Ice Stream: Robert Bell on Hudson Strait. *Géographie Physique et Quaternaire* **61**(2–3): 211–215.
- Canals M, Urgeles R, Calafat AM. 2000. Deep sea-floor evidence of past ice streams off the Antarctic Peninsula. *Geology* **28**(1): 31–34.
- Carr JR, Stokes CR, Vieli A. 2013. Recent progress in understanding marine-terminating Arctic outlet glacier response to climatic and oceanic forcing: twenty years of rapid change. *Progress in Physical Geography* **37**(4): 435–466.
- Carsey F, Behar A, Lane AL, Realmuto V, Engelhardt H. 2002. A borehole camera system for imaging the deep interior of ice sheets. *Journal of Glaciology* **48**(163): 622–628.
- Carter SP, Fricker HA. 2012. The supply of meltwater to the grounding line of the Siple coast, West Antarctica. *Annals of Glaciology* **53**(6): 267–280.
- Carter SP, Fricker HA, Siegfried MR. 2017. Antarctic subglacial lakes drain through sediment-floored canals: theory and model testing on real and idealised domains. *The Cryosphere* **11**: 381–405.
- Chapwanya M, Clark CD, Fowler AC. 2011. Numerical computations of a theoretical model of ribbed moraine formation. *Earth Surface Processes and Landforms* **36**: 1105–1112.
- Christofferson P, Tulaczyk S. 2003a. Thermodynamics of basal freeze-on: predicting basal and subglacial signatures of stopped ice streams and interstream ridges. *Annals of Glaciology* **36**: 233–243.
- Christofferson P, Tulaczyk S. 2003b. Response of subglacial sediments to basal freeze-on: 1. Theory and comparison to observations from beneath the west Antarctic Ice Sheet. *Journal of Geophysical Research* **108**(B4): 19.1–19.16.
- Christofferson P, Tulaczyk S, Behar A. 2010. Basal ice sequences in Antarctic ice stream: exposure of past hydrologic conditions and a principal mode of sediment transfer. *Journal of Geophysical Research* **115**: F03034, <https://doi.org/10.1029/2009JF001430>.
- Clark CD. 1993. Mega-scale glacial lineations and cross-cutting ice flow landforms. *Earth Surface Processes and Landforms* **18**(1): 1–29.
- Clark CD. 2010. Emergent drumlins and their clones: from till dilatancy to flow instabilities. *Journal of Glaciology* **51**(200): 1011–1025.
- Clark CD, Hughes ALC, Greenwood SL, Spagnolo M, Ng FSL. 2009. Size and shape characteristics of drumlins, derived from a large sample, and associated scaling laws. *Quaternary Science Reviews* **28**(7–8): 677–692.
- Clark CD, Stokes CR. 2001. Extent and basal characteristics of the M'Clintock Channel Ice Stream. *Quaternary International* **86**: 81–101.
- Clark CD, Tulaczyk SM, Stokes CR, Canals M. 2003. A groove-ploughing theory for the production of mega-scale glacial lineations, and implications for ice-stream mechanics. *Journal of Glaciology* **49**: 240–256.
- Clarke GKC. 1987. Fast glacier flow: ice streams, surging, tidewater glaciers. *Journal of Geophysical Research* **92**(B9): 8835–8841.
- Clarke GKC, Leverington DW, Telle JT, Dyke AS, Marshall SJ. 2005. Fresh arguments against the Shaw megaflood hypothesis. A reply to comments by David Sharpe on 'Palaeohydraulics of the last outburst flood from glacial-Lake Agassiz and the 8200 BP cold event'. *Quaternary Science Reviews* **24**: 1533–1541.
- Clarke TS, Echelmeyer K. 1996. Seismic-reflection evidence for a deep subglacial trough beneath Jakobshavn Isbræ, West Greenland. *Journal of Glaciology* **42**(141): 219–232.
- Clayton L, Teller JT, Attig JW, Jr. 1985. Surging of the southwestern part of the Laurentide Ice Sheet. *Boreas* **14**(3): 235–241.
- Conway H, Hall BL, Denton GH, Gades AM, Waddington ED. 1999. Past and future grounding-line retreat of the West Antarctic Ice Sheet. *Science* **286**: 280–283.
- Cook SJ, Swift D. 2012. Subglacial basins: their origin and importance in glacial systems and landscapes. *Earth-Science Reviews* **115**(4): 332–372.
- Cowton T, Nienow P, Bartholomew I, Sole A, Mair D. 2012. Rapid erosion beneath the Greenland Ice Sheet. *Geology* **40**(4): 343–346.
- Crabtree RD, Doake CSM. 1982. Pine Island Glacier and its drainage basin: results from radio-echo sounding. *Annals of Glaciology* **3**: 65–70.
- Cuffrey K, Alley RB. 1996. Is erosion by deforming subglacial sediments significant? (towards till continuity). *Annals of Glaciology* **22**: 17–24.
- De Angelis H, Kleman J. 2008. Palaeo-ice stream onset: examples from the north-eastern Laurentide Ice Sheet. *Earth Surface Processes and Landforms* **33**(4): 560–572.
- Dean WG. 1953. The drumlinoid land forms of the Barren Grounds. *Canadian Geographer* **1**: 19–30.
- Denton GH, Hughes T. 1981. *The Last Great Ice Sheets*. Wiley Interscience: New York; 484.
- Doake CSM, Corr HFJ, Jenkins A, Makinson K, Nicholls KW, Nath C, Smith AM, Vaughan DG. 2001. Rutford Ice Stream, West Antarctica. In *The West Antarctic Ice Sheet: Behaviour and Environment*, Alley

- RB, Bindschadler RA (eds), Vol. 77, American Geophysical Union: Antarctic Research Series; 221–235.
- Doake CSM, Frolich RM, Mantripp DR, Smith AM, Vaughan DG. 1987. Glaciological studies on Rutford Ice Stream, Antarctica. *Journal of Geophysical Research - Earth Surface* **92**(B9): 8951–8960.
- Domack EW, Jacobson EA, Shipp SS, Anderson JB. 1999. Late Pleistocene–Holocene retreat of the West Antarctic Ice-Sheet system in the Ross Sea: Part 2 – Sedimentologic and stratigraphic signature. *Geological Society of America Bulletin* **111**: 1517–1536.
- Dowdeswell JA, Canals M, Jakobsson M, Todd BJ, Dowdeswell EK, Hogan K (eds). 2016. Atlas of Submarine Glacial Landforms: Modern, Quaternary and Ancient. *Geological Society Memoirs* **46**: 618.
- Dowdeswell JA, Fugelli EMG. 2012. The seismic architecture and geometry of grounding-zone wedges formed at the marine margins of past ice sheets. *Geological Society of America Bulletin* **124**(11–12): 1750–1761.
- Dowdeswell JA, Kenyon NH, Elverhøi A, Laberg JS, Hollender F-J, Mienert J, Siegert MJ. 1996. Large-scale sedimentation on the glacier-influenced Polar North Atlantic margins: long-range side-scan sonar evidence. *Geophysical Research Letters* **23**: 3535–3538.
- Dowdeswell JA, Maslin MA, Andrews JT, McCave IN. 1995. Iceberg production, debris rafting, and the extent and thickness of Heinrich layers (H-1, H-2) in North Atlantic sediments. *Geology* **23**: 301–304.
- Dowdeswell JA, Ó Cofaigh C, Pudsey CJ. 2004. Thickness and extent of the subglacial till layer beneath an Antarctic palaeo-ice stream. *Geology* **32**: 13–16.
- Dowdeswell JA, Ottesen D, Evans J, Ó Cofaigh C, Anderson JB. 2008. Submarine glacial landforms and rates of ice-stream collapse. *Geology* **36**: 819–822.
- Dredge LA, Cowan WR. 1989. Quaternary geology of the southwestern Canadian Shield. In *Quaternary Geology of Canada and Greenland*, Fulton RJ (ed). Geological Survey of Canada: Ottawa; 214–235.
- Dunlop P, Clark CD. 2006. The morphological characteristics of ribbed moraine. *Quaternary Science Reviews* **25**: 1668–1691.
- Dunlop P, Clark CD, Hindmarsh RCA. 2008. Bed ribbing instability explanation: testing a numerical model of ribbed moraine formation arising from coupled flow of ice and subglacial sediment. *Journal of Geophysical Research* **113**: F03005.
- Dyke AS. 1984. Quaternary geology of Boothia Peninsula and northern District of Keewatin, central Canadian Arctic. *Geological Survey of Canada Memoir* **407**: 26.
- Dyke AS, Morris TF. 1988. Canadian landform examples. 7. Drumlin fields, dispersal trins, and ice streams in Arctic Canada. *Canadian Geographer* **32**(1): 86–90.
- Dyke AS, Morris TF, Green DEC, England J. 1992. Quaternary geology of Prince of Wales Island, Arctic Canada. *Geological Survey of Canada Memoir* **433**: 142.
- Dyke AS, Prest VK. 1987. Late Wisconsinan and Holocene history of the Laurentide Ice Sheet. *Géographie Physique et Quaternaire* **41**(2): 237–263.
- Echelmeyer KA, Harrison WD, Larsen C, Mitchell JE. 1994. The role of the margins in the dynamics of an active ice stream. *Journal of Glaciology* **40**(136): 527–538.
- Edgeworth TW. 1914. Antarctica and some of its problems. *The Geographical Journal* **43**(6): 605–627.
- Elverhøi A, Norem H, Anderson ES, Dowdeswell JA, Fossen I, Hafliðason H, Kenyon NH, Laberg JS, King EL, Sejrup HP, Solheim A, Vorren T. 1997. On the origin and flow behaviour of submarine slides on deep-sea fans along the Norwegian-Barents Sea continental margin. *Geo-Marine Letters* **17**: 119–125.
- Ely JC, Clark CD, Spagnolo M, Stokes CR, Greenwood SL, Hughes ALC, Dunlop P, Hess D. 2016. Do subglacial bedforms comprise a size and shape continuum? *Geomorphology* **257**: 108–119.
- Engelhardt H, Humphrey N, Kamb B, Fahnestock M. 1990. Physical conditions at the base of a fast moving Antarctica ice stream. *Science* **248**(4951): 57–59.
- Engelhardt H, Kamb B. 1998. Basal sliding of Ice Stream B, West Antarctica. *Journal of Glaciology* **44**(147): 223–230.
- Evans DJA, Clark CD, Rea BR. 2008. Landform and sediment imprints of fast glacier flow in the southwest Laurentide ice sheet. *Journal of Quaternary Science* **46**: 80–125.
- Evans DJA, Lemmen DS, Rea BR. 1999. Glacial landforms of the southwest Laurentide ice sheet: modern Icelandic analogues. *Journal of Quaternary Science* **14**(7): 673–691.
- Evans DJA, Young NJP, Ó Cofaigh C. 2014. Glacial geomorphology of the terrestrially-terminating fast flow lobes/ice stream margins in the southwest Laurentide ice sheet. *Geomorphology* **204**: 86–113.
- Evans J, Ó Cofaigh C, Dowdeswell JA, Wadhams P. 2009. Marine geophysical evidence for former expansion and flow of the Greenland Ice Sheet across the northeast Greenland continental shelf. *Journal of Quaternary Science* **24**: 279–293.
- Eyles N. 2012. Rock drumlins and megaflutes of the Niagara Escarpment, Ontario, Canada: a hard bed landform assemblage cut by the Saginaw-Huron Ice Stream. *Quaternary Science Reviews* **55**: 34–49.
- Favier L, Durand G, Cornford SL, Gudmundsson GH, Gagliardini O, Gillet-chaulet F, Zwinger T, Payne AJ, Le Brocq AM. 2014. Retreat of Pine Island Glacier controlled by marine ice-sheet stability. *Nature Climate Change* **4**: 117–121.
- Fisher TG, Shaw J. 1992. A depositional model for Rogen moraine, with examples from the Avalon Peninsula, Newfoundland. *Canadian Journal of Earth Sciences* **29**: 669–686.
- Fowler AC. 2003. On the rheology of till. *Annals of Glaciology* **37**: 55–59.
- Fowler AC. 2010. The formation of subglacial streams and mega-scale glacial lineations. *Proceedings of the Royal Society A* **466**: 3181–3201.
- Fowler AC, Chapwanya M. 2014. An instability theory for the formation of ribbed moraine, drumlins and mega-scale glacial lineations. *Proceedings of the Royal Society A* **470**: 20140185.
- Fowler AC, Spagnolo M, Clark CD, Stokes CR, Hughes ALC, Dunlop P. 2013. On the size and shape of drumlins. *International Journal on Geomathematics* **4**: 155–165.
- Fricter HA, Scambos T. 2009. Connected subglacial lake activity on lower Mercer and Whillans Ice Streams, West Antarctica, 2003–2008. *Journal of Glaciology* **55**(190): 303–315.
- Fricter HA, Scambos T, Bindschadler R, Padman L. 2007. An active subglacial water system in West Antarctica mapped from space. *Science* **315**(5818): 1544–1548.
- Fricter HA, Scambos T, Carter S, Davis C, Haran T, Joughin I. 2010. Synthesizing multiple remote-sensing techniques for subglacial hydrologic mapping: application to a lake system beneath MacAyeal Ice Stream, West Antarctica. *Journal of Glaciology* **56**(196): 187–199.
- Fricter HA, Siegfried MR, Carter SP, Scambos TA. 2016. A decade of progress in observing and modelling Antarctic subglacial water systems. *Philosophical Transactions of the Royal Society A* **374**: 20140294.
- Frolich RM, Doake CSM. 1988. Relative importance of lateral and vertical shear on Rutford Ice Stream, Antarctica. *Annals of Glaciology* **11**: 19–22.
- Graham AGC, Dutrieux P, Vaughan DG, Nitsche FO, Gyllencreutz R, Greenwood SL, Larter RD, Jenkins A. 2013. Seabed corrugations beneath an Antarctica ice shelf revealed by autonomous underwater vehicle survey: origin and implications for the history of Pine Island Glacier. *Journal of Geophysical Research* **118**: 1–11. <https://doi.org/10.1002/jgrf.20087>.
- Graham AGC, Larter RD, Gohl K, Dowdeswell JA, Hillenbrand C-D, Smith JA, Evans J, Kuhn G, Deen T. 2010. Flow and retreat of the Late Quaternary Pine Island-Thwaites palaeo-ice stream, West Antarctica. *Journal of Geophysical Research* **115**: F03025.
- Graham AGC, Larter RD, Gohl K, Hillenbrand C-D, Smith JA, Kuhn G. 2009. Bedform signature of a West Antarctic palaeo-ice stream reveals a multi-temporal record of flow and substrate control. *Quaternary Science Reviews* **28**: 2774–2793.
- Gravenor CP, Meneley WA. 1958. Glacial flutings in central and northern Alberta. *American Journal of Science* **256**: 715–728.
- Gray L, Joughin I, Tulaczyk S, Spikes VB, Bindschadler R, Jezek K. 2005. Evidence for subglacial water transport in the West Antarctic Ice Sheet through three-dimensional satellite radar interferometry. *Geophysical Research Letters* **32**: L03501. <https://doi.org/10.1029/2004GL021387>.
- Greenwood SL, Kleman J. 2010. Glacial landforms of extreme size in the Keewatin sector of the Laurentide ice sheet. *Quaternary Science Reviews* **29**: 1894–1910.

- Hallet B, Hunter L, Bogen J. 1996. Rates of erosion and sediment evacuation by glaciers: a review of field data and their implications. *Global and Planetary Change* **12**: 213–235.
- Hanna E, Navarro FJ, Pattyn F, Domingues CM, Fettweis X, Ivins ER, Nicholls RJ, Ritz C, Smith B, Tulaczyk S, Whitehouse PL, Zwally HJ. 2013. Ice sheet mass balance and climate change. *Nature* **498**: 51–59.
- Hart JK. 1997. The relationship between drumlins and other forms of subglacial glaciotectionic deformation. *Quaternary Science Reviews* **16**: 93–107.
- Hart JK. 1999. Identifying fast ice flow from landform assemblages in the geological record: a discussion. *Annals of Glaciology* **28**: 59–66.
- Hättestrand C. 1997. Ribbed moraines in Sweden – distribution, pattern and palaeoglaciological implications. *Sedimentary Geology* **111**: 41–56.
- Hättestrand C, Kleman J. 1999. Ribbed moraine formation. *Quaternary Science Reviews* **18**: 43–61.
- Hicock SR. 1988. Calcareous till facies north of Lake Superior, Ontario: implications for Laurentide ice streaming. *Géographie Physique et Quaternaire* **42**(2): 120–135.
- Hicock SR, Dreimanis A. 1992. Deformation till in the Great Lakes region: implications for rapid flow along the south-central margin of the Laurentide Ice Sheet. *Canadian Journal of Earth Sciences* **29**(7): 1565–1579.
- Hillier JK, Kougioumtzoglou IA, Stokes CR, Smith MJ, Clark CD, Spagnolo M. 2016. Exploring explanations of subglacial bedform sizes using statistical models. *PLoS One* **11**(7): e0159489.
- Hillier JK, Smith MJ, Clark CD, Stokes CR, Spagnolo M. 2013. Subglacial bedforms reveal an exponential size-frequency distribution. *Geomorphology* **190**: 82–91.
- Hindmarsh RCA. 1997. Deforming beds: viscous and plastic scales of deformation. *Quaternary Science Reviews* **16**(9): 1039–1056.
- Hindmarsh RCA. 1998a. The stability of a viscous till sheet coupled with ice flow, considered at wavelengths less than the ice thickness. *Journal of Glaciology* **44**(147): 285–292.
- Hindmarsh RCA. 1998b. Drumlinization and drumlin-forming instabilities: viscous till mechanisms. *Journal of Glaciology* **44**(147): 293–314.
- Hindmarsh RCA, Stokes CR. 2008. Formation mechanisms for ice-stream lateral shear margin moraines. *Earth Surface Processes and Landforms* **33**(4): 610–626.
- Hodgson DA. 1993. Surficial Geology, Storkerson Peninsula, Victoria Island and Stefansson Island, Northwest Territories. *Geological Survey of Canada, Map 1817A, Scale 1:250 000*.
- Hodgson DA. 1994. Episodic ice streams and ice shelves during retreat of the northwesternmost sector of the late Wisconsinan Laurentide Ice Sheet over the central Canadian Arctic Archipelago. *Boreas* **23**(1): 14–28.
- Hooke RL, Elverhøi A. 1996. Sediment flux from a fjord during glacial periods, Isfjorden, Spitsbergen. *Global and Planetary Change* **12**: 237–249.
- Horgan HJ, Anandakrishnan S, Jacobel RW, Christianson K, Alley RB, Heeszel DS, Picotti S, Walter JJ. 2012. Subglacial Lake Whillans – seismic observations of a shallow active reservoir beneath a west Antarctic ice stream. *Earth and Planetary Science Letters* **331–332**: 201–209.
- Hughes TJ. 1972. *Is the West Antarctic Ice Sheet disintegrating?* ISCAP [Ice Streamline Cooperative Antarctic Project] Bulletin. Ohio State University, No. 1.
- Hughes TJ. 1973. Is the West Antarctic Ice Sheet disintegrating? *Journal of Geophysical Research* **78**: 7884–7910.
- Hughes TJ. 1977. West Antarctic ice streams. *Reviews of Geophysics and Space Physics* **15**(1): 1–46.
- Humphrey NF, Kamb B, Fahnestock M, Engelhardt H. 1993. Characteristics of the bed of the lower Columbia Glacier, Alaska. *Journal of Geophysical Research* **98**: 837–846.
- Humphrey NF, Raymond CF. 1994. Hydrology, erosion and sediment production in a surging glacier: Variegated Glacier, Alaska, 1982–1983. *Journal of Glaciology* **40**: 539–552.
- Iken A, Echelmeyer K, Harrison W, Funk M. 1993. Mechanisms of fast flow in Jakobshavn Isbræ, West Greenland: Part I. Measurements of temperature and water level in deep boreholes. *Journal of Glaciology* **39**(131): 15–25.
- Iverson NR. 2010. Shear resistance and continuity of subglacial till: hydrology rules. *Journal of Glaciology* **56**(200): 1104–1115.
- Iverson NR, Bwaker RW, Hooyer TS. 1997. A ring-shear device for the study of till deformation: tests on tills with contrasting clay contents. *Quaternary Science Reviews* **16**(9): 1057–1066.
- Iverson NR, Hooyer TS, Baker RW. 1998. Ring-shear studies of till deformation: Coulomb-plastic behaviour and distributed strain in glacier beds. *Journal of Glaciology* **44**(148): 634–642.
- Iverson NR, Janson B, LeB Hooke R, Jansson P. 1995. Flow mechanism of glaciers on soft beds. *Science* **267**(5194): 80–81.
- Iverson NR, Jansson P, LeB Hooke R. 1994. *In situ* measurement of the strength of deforming subglacial till. *Journal of Glaciology* **40**(136): 497–503.
- Jakobsson M, Anderson JB, Nitsche FO, Dowdeswell JA, Gyllencreutz R, Kirchner N, Mohammad R, O'Regan M, Alley RB, Anandakrishnan S, Eriksson B, Kirshner A, Fernandez R, Stollendorf T, Minzoni R, Majewski W. 2011. Geological record of ice shelf break-up and grounding line retreat, Pine Island Bay, West Antarctica. *Geology* **39**: 691–694.
- Jamieson SR, Vieli A, Livingstone SJ, Ó Cofaigh C, Stokes CR, Hillenbrand CD, Dowdeswell JA. 2012. Ice-stream stability on a reverse bed slope. *National Geoscience* **5**: 799–802.
- Jenson JW, Clark PU, MacAyeal DR, Ho C, Vela JC. 1995. Numerical modelling of advective transport of saturated deforming sediment beneath the Lake Michigan Lobe, Laurentide Ice Sheet. *Geomorphology* **14**: 157–166.
- Jezek K, Wu X, Gogineni P, Rodríguez E, Freeman A, Rodríguez-Morales F, Clark CD. 2011. Radar images of the bed of the Greenland Ice Sheet. *Geophysical Research Letters* **38**: L01501. <https://doi.org/10.1029/2010GL045519>.
- Joughin I, Gray L, Bindschadler R, Price S, Morse D, Hulbe C, Mattar K, Werner C. 1999. Tributaries of West Antarctic ice streams revealed by RADARSAT interferometry. *Science* **286**: 283–286.
- Joughin I, MacAyeal DR, Tulaczyk S. 2004. Basal shear stress of the Ross ice streams from control method inversions. *Journal of Geophysical Research* **109**: B09405. <https://doi.org/10.1029/2003JB002960>.
- Joughin IR, Smith BE, Medley B. 2014. Marine ice sheet collapse potentially under way for the Thwaites Glacier Basin, West Antarctica. *Science* **344**: 735–738.
- Joughin IR, Tulaczyk S, Bamber JL, Blankenship DD, Holt JW, Scambos T, Vaughan DG. 2009. Basal conditions for Pine Island and Thwaites Glaciers, West Antarctica, determined using satellite and airborne data. *Journal of Glaciology* **55**(190): 245–257.
- Kamb B. 1991. Rheological nonlinearity and flow instability in the deforming bed mechanism of ice stream motion. *Journal of Geophysical Research* **96**(B10): 16 585–16 595 In *The West Antarctic Ice Sheet: Behaviour and Environment* Alley RB, Bindschadler RA (eds). Antarctic Research Series, vol. 77. American Geophysical Union, Washington DC, 105–121.
- Kamb B. 2001. Basal zone of the West Antarctic ice streams and its role in lubrication of their rapid motion. In Alley RB, Bindschadler RA (Eds) *The West Antarctic Ice Sheet: Behaviour and Environment*, Alley RB, Bindschadler RA (eds). American Geophysical Union, Antarctic Research Series **77**: 157–199.
- King EC, Hindmarsh RCA, Stokes CR. 2009. Formation of mega-scale glacial lineations observed beneath a West Antarctic ice stream. *Nature Geoscience* **2**: 585–588.
- King E, Pritchard H, Brisbourne A, Smith A, Lewington E, Clark C. 2017. Comparison of contemporary subglacial bedforms beneath Rutford and Pine Island Glacier, Antarctica. Abstract from the 19th EGU General Assembly, EGU 2017, proceedings from the conference held 23–28 April, 2017 in Vienna, Austria, p. 1885.
- King EC, Pritchard HD, Smith AM. 2016. Subglacial landforms beneath Rutford Ice Stream, Antarctica: detailed bed topography from ice-penetrating radar. *Earth System Science Data* **8**: 151–158.
- King EC, Woodward J, Smith AM. 2004. Seismic evidence for a water-filled canal in deforming till beneath Rutford Ice Stream, West Antarctica. *Geophysical Research Letters* **31**: L20401. <https://doi.org/10.1029/2004GL020379>.
- King EC, Woodward J, Smith AM. 2007. Seismic and radar observations of subglacial bed forms beneath the onset zone of Rutford Ice Stream Antarctica. *Journal of Glaciology* **53**: 665–672.

- King EL, Sejrup HP, Haflidason H, Elverhøi A, Aarseth I. 1996. Quaternary seismic stratigraphy of the North Sea Fan: glacial-fed gravity flow aprons, hemipelagic sediments, and large submarine slides. *Marine Geology* **130**: 296–315.
- Kite ES, Hindmarsh RCA. 2007. Did ice streams shape the largest channels on Mars? *Geophysical Research Letters* **34**(19): L19202.
- Kleman J, Hättestrand C. 1999. Frozen-bed Fennoscandian and Laurentide ice sheets during the Last Glacial Maximum. *Nature* **402**: 63–66.
- Kleman J, Borgström I. 1996. Reconstruction of palaeo-ice sheets: the use of geomorphological data. *Earth Surface Processes and Landforms* **21**(10): 893–909.
- Knight J. 2002. Glacial sedimentary evidence supporting stick-slip basal ice flow. *Quaternary Science Reviews* **21**: 975–983.
- Krabbendam M, Eyles N, Putkinen N, Bradwell T, Arbelaez-Moreno L. 2016. Streamlined hard beds formed by palaeo-ice streams: a review. *Sedimentary Geology* **338**: 24–50.
- Laberg JS, Vorren TO. 1995. Late Weichselian submarine debris flow deposits on the Bear Island Trough Mouth Fan. *Marine Geology* **127**: 45–72.
- Larter RD, Graham AGC, Gohl K, Kuhn G, Hillenbrand C-D, Smith JA, Deen TJ, Livermore RA, Schenke H-W. 2009. Subglacial bedforms reveal complex basal regime in a zone of paleo-ice stream convergence, Amundsen Sea embayment, West Antarctica. *Geology* **37**(5): 411–414.
- Le Brocq AM, Payne AJ, Siegert MJ, Alley RB. 2009. A subglacial water-flow model for West Antarctica. *Journal of Glaciology* **55**: 879–888.
- Le Brocq AM, Ross N, Griggs JA, Bingham RG, Corr HFJ, Ferraccioli F, Jenkins A, Jordan TA, Payne AJ, Rippin DM, Siegert MJ. 2013. Evidence from ice shelves for channelized meltwater flow beneath the Antarctic Ice Sheet. *Nature Geoscience* **6**: 945–948.
- Lian O, Hicock SR, Dreimanis A. 2003. Laurentide and Cordilleran fast ice flow: some sedimentological evidence from Wisconsinan subglacial till and its substrate. *Boreas* **32**(1): 102–113.
- Lien R, Solheim A, Elverhøi A, Rokoengen K. 1989. Iceberg scouring and sea bed morphology on the eastern Weddell Sea shelf, Antarctica. *Polar Research* **7**(1): 43–57.
- Lingle CS, Brown TJ. 1987. A subglacial aquifer bed model and water pressure dependent basal sliding relationship for a West Antarctic ice stream. In *Dynamics of the West Antarctic Ice Sheet*, van der Veen CJ, Oerlemans J (eds), Proceedings of a workshop held in Utrecht, May 6–8, 1985. D. Reidel Publishing Company: Dordrecht; 249–285.
- Livingstone SJ, Ó Cofaigh CO, Stokes CR, Hillenbrand C-D, Vieli A, Jamieson SR. 2012. Antarctic palaeo-ice streams. *Earth-Science Reviews* **111**: 90–128.
- Livingstone SJ, Stokes CR, Ó Cofaigh C, Hillenbrand C-D, Vieli A, Jamieson SSR, Spagnolo M, Dowdeswell JA. 2016a. Subglacial processes on an Antarctic ice stream bed. 1: sediment transport and bedform genesis inferred from marine geophysical data. *Journal of Glaciology* **62**(232): 270–284.
- Livingstone SJ, Utting DJ, Ruffell A, Clark CD, Pawley S, Atkinson N, Fowler AC. 2016b. Discovery of relict subglacial lakes and their geometry and mechanism of drainage. *Nature Communications* **7**: 11767. <https://doi.org/10.1038/ncomms11767>.
- Lliboutry L. 1987. Realistic, yet simple bottom-boundary conditions beneath glaciers and ice sheets. *Journal of Geophysical Research* **92**(B9): 9101–9109.
- Luchitta BK, Anderson DM, Shoji H. 1981. Did ice streams carve Martian outflow channels? *Nature* **290**: 759–763.
- Luternauer JL, Murray JW. 1983. Late Quaternary morphologic development and sedimentation, central British Columbia continental shelf. *Geological Survey of Canada Paper* **83-21**: 38.
- MacAyeal DR. 1992. The basal stress distribution of Ice Stream E, Antarctica, inferred by control methods. *Journal of Geophysical Research* **97**(B1): 9101–9109.
- MacAyeal DR, Bindschadler RA, Scambos TA. 1995. Basal friction of Ice Stream E, west Antarctica. *Journal of Glaciology* **5**(41): 661–603.
- MacGregor JA, Catania GA, Conway H, Schroeder DM, Joughin I, Young DA, Kempf SD, Blankenship DD. 2013. Weak bed control of the eastern shear margin of Thwaites Glacier, West Antarctica. *Journal of Glaciology* **59**(217): 900–912.
- Margold M, Stokes CR, Clark CD. 2015a. Ice streams in the Laurentide Ice Sheet: identification, characteristics and comparison to modern ice sheets. *Earth-Science Reviews* **143**: 117–146.
- Margold M, Stokes CR, Clark CD, Kleman J. 2015b. Ice streams of the Laurentide Ice Sheet: a new mapping inventory. *Journal of Maps* **11**(3): 380–395.
- Mathews WH. 1974. Surface profiles of the Laurentide Ice Sheet in its marginal areas. *Journal of Glaciology* **13**: 37–43.
- Mathews WH. 1991. Ice sheets and ice streams: thoughts on the Cordilleran Ice Sheet symposium. *Géographie Physique et Quaternaire* **45**(3): 263–267.
- McIntyre NF. 1985. The dynamics of ice sheet outlets. *Journal of Glaciology* **31**(108): 99–107.
- Meier MF, Post A. 1969. What are glacier surges? *Canadian Journal of Earth Sciences* **6**(4): 807–817.
- Mercer JH. 1978. West Antarctic ice sheet and CO<sub>2</sub> greenhouse effect: a threat of disaster. *Nature* **271**: 321–325.
- Morgan VI, Budd WF. 1975. Radio-echo sounding of the Lambert Glacier basin. *Journal of Glaciology* **15**(73): 103–111.
- Morlighem M, Rignot E, Mouginot J, Seroussi H, Larour E. 2014. Deeply incised submarine glacial valleys beneath the Greenland Ice Sheet. *Nature Geoscience* **7**: 418–422.
- Mosola AB, Anderson JB. 2006. Expansion and rapid retreat of the West Antarctic Ice Sheet in eastern Ross Sea: possible consequence of over-extended ice streams? *Quaternary Science Reviews* **25**: 2177–2196.
- Murray T. 1997. Assessing the paradigm shift: deformable glacier beds. *Quaternary Science Reviews* **16**: 995–1016.
- Murray T, Corr H, Forieri A, Smith AM. 2008. Contrasts in hydrology between regions of basal deformation and sliding beneath Rutford ice stream, West Antarctica, mapped using radar and seismic data. *Geophysical Research Letters* **35**. <https://doi.org/10.1029/2008GL033681>.
- Ng FSL. 2000. Canals under sediment-based ice sheets. *Annals of Glaciology* **30**: 146–152.
- Nick FM, Vieli A, Anderson ML, Joughin I, Payne A, Edwards TL, Pattyn F, van de Wal RSW. 2013. Future sea-level rise from Greenland's main outlet glaciers in a warming climate. *Nature* **497**: 235–238.
- Novick AN, Bentley CR, Lord N. 1994. Ice thickness, bed topography and basal-reflection strengths from radar sounding, Upstream B, West Antarctica. *Annals of Glaciology* **20**: 148–152.
- Nygård A, Sejrup HP, Haflidason H, Lekens WAH, Clark CD, Bigg GR. 2007. Extreme sediment and ice discharge from marine-based ice streams: new evidence from the North Sea. *Geology* **35**(5): 395–398.
- Ó Cofaigh C. 2012. Ice sheets viewed from the ocean: the contribution of marine science to understanding modern and past ice sheets. *Philosophical Transactions of the Royal Society A* **370**: 5512–5539.
- Ó Cofaigh C, Dowdeswell JA, Allen CS, Hiemstra J, Pudsey CJ, Evans J, Evans DJA. 2005. Flow dynamics and till genesis associated with a marine-based Antarctic palaeo-ice stream. *Quaternary Science Reviews* **24**: 709–740.
- Ó Cofaigh C, Dowdeswell JA, Evans J, Larter RD. 2008. Geological constraints on Antarctic palaeo-ice-stream retreat. *Earth Surface Processes and Landforms* **33**: 513–525.
- Ó Cofaigh C, Dowdeswell JD, King EC, Anderson JB, Clark CD, Evans DJA, Evans J, Hindmarsh RCA, Larter RD, Stokes CR. 2010a. Comment on Shaw J, Pugin A and Young R (2008): 'A meltwater origin for Antarctic shelf bedforms with special attention to megalineations'. *Geomorphology* **102**: 364–375 *Geomorphology* **117**: 195–198.
- Ó Cofaigh C, Evans DJA, Smith IR. 2010b. Large-scale reorganisation and sedimentation of terrestrial ice streams during late Wisconsinan Laurentide ice sheet deglaciation. *Geological Society of America Bulletin* **122**(5–6): 743–756.
- Ó Cofaigh C, Evans J, Dowdeswell JA, Larter RD. 2007, 112. Till characteristics, genesis and transport beneath Antarctic paleo-ice streams. *Journal of Geophysical Research* F03006. <https://doi.org/10.1029/2006JF000606>.
- Ó Cofaigh C, Pudsey CJ, Dowdeswell JA, Morris P. 2002. Evolution of subglacial bedforms along a paleo-ice stream, Antarctic Peninsula continental shelf. *Geophysical Research Letters* **29**(8): 1199.

- Ó Cofaigh C, Stokes CR, Lian OB, Clark CD, Tulaczyk SM. 2013. Formation of mega-scale glacial lineations on the Dubawnt Lake Ice Stream bed: 2. Sedimentology and stratigraphy. *Quaternary Science Reviews* **77**: 190–209.
- Ottesen D, Dowdeswell JA. 2006. Assemblages of submarine landforms produced by tidewater glaciers in Svalbard. *Journal of Geophysical Research* **111**: F01016.
- Ottesen D, Dowdeswell JA. 2009. An inter-ice stream glaciated margin: submarine landforms and a geomorphic model based on marine-geophysical data from Svalbard. *Geological Society of America Bulletin* **121**: 1647–1665.
- Ottesen D, Dowdeswell JA, Rise L. 2005. Submarine landforms and the reconstruction of fast-flowing ice streams within a large Quaternary ice sheet: the 2500-km-long Norwegian-Svalbard margin (57°–80°N). *Geological Society of America Bulletin* **117**(7/8): 1033–1050.
- Ottesen D, Stokes CR, Rise L, Olsen L. 2008. Ice-sheet dynamics and ice streaming along the coastal parts of northern Norway. *Quaternary Science Reviews* **27**: 922–940.
- Paterson WSB. 1994. *The Physics of Glaciers* (3rd edn). Pergamon: Oxford; 480.
- Patterson CJ. 1997. Southern Laurentide ice lobes were created by ice streams: des Moines lobe in Minnesota, USA. *Sedimentary Geology* **111**(1–4): 249–261.
- Patterson CJ. 1998. Laurentide glacial landscapes: the role of ice streams. *Geology* **26**(7): 643–646.
- Payne AJ, Dongelmans PW. 1997. Self-organisation on the thermo-mechanical flow of ice sheets. *Journal of Geophysical Research* **102**: 12219–12234.
- Peters LE, Anandakrishnan S, Alley RB, Smith AM. 2007. Extensive storage of basal meltwater in the onset region of a major West Antarctic ice stream. *Geology* **35**(3): 251–254.
- Pfeffer WT, Harper JT, O'Neel S. 2008. Kinematic constraints on glacier contributions to 21st Century sea level rise. *Science* **321**: 1340–1343.
- Piotrowski JA, Larsen NK, Junge FW. 2004. Reflections on soft subglacial beds as a mosaic of deforming and stable spots. *Quaternary Science Reviews* **23**: 993–1000.
- Price SF, Bindschadler RA, Hulbe CL, Blankenship DD. 2002. Force balance along an inland tributary and onset to Ice Stream D, West Antarctica. *Journal of Glaciology* **48**(160): 20–30.
- Price SF, Bindschadler RA, Hulbe CL, Joughin IR. 2001. Post-stagnation behaviour in the upstream regions of Ice Stream C, West Antarctica. *Journal of Glaciology* **47**(157): 283–294.
- Punkari M. 1995a. Function of the ice streams in the Scandinavian Ice Sheet: analyses of glacial geological data from southwestern Finland. *Transactions of the Royal Society of Edinburgh: Earth Sciences* **85**: 283–302.
- Punkari M. 1995b. Glacial flow systems in the zone of confluence between the Scandinavian and Novaya Zemlya Ice Sheets. *Quaternary Science Reviews* **14**: 589–603.
- Punkari M. 1997. Glacial and glaciofluvial deposits in the interlobate areas of the Scandinavian Ice Sheet. *Quaternary Science Reviews* **16**: 741–753.
- Raymond CF, Echelmeyer KA, Whillans IM, Doakes CSM. 2001. Ice stream shear margins. In *The West Antarctic Ice Sheet: Behaviour and Environment*, Alley RB, Bindschadler RA (eds), Vol. 77, Antarctic Research Series. American Geophysical Union: Washington DC; 137–155.
- Rink H. 1877. *Danish Greenland: its People and Products*. H.S. King: London; 648.
- Rippin DM, Bamber JL, Siegert MJ, Vaughan DG, Corr HFJ. 2006. Basal conditions beneath enhanced-flow tributaries of Slessor Glacier, East Antarctica. *Journal of Glaciology* **52**: 481–490.
- Ritz C, Edwards TL, Durand G, Payne AJ, Peyaud V, Hindmarsh RCA. 2015. Potential sea-level rise from Antarctic ice-sheet instability constrained by observations. *Nature* **528**: 115–118.
- Robin G de Q, Evans S, Drewry DJ, Harrison CH, Petrie DL. 1970. Radio-echo sounding of the Antarctic Ice Sheet. *Antarctic Journal of the United States* **5**(6): 229–232.
- Rooney ST, Blankenship DD, Alley RB, Bentley CR. 1987a. Till beneath Ice Stream B 2. Structure and continuity. *Journal of Geophysical Research* **92**(B9): 8913–8920.
- Rooney ST, Blankenship DD, Bentley CR. 1987b. Seismic refraction measurements of crustal structure in West Antarctica. In *Gondwana Six: Structure, Tectonics and Geophysics*, McKenzie GD (ed), Vol. 40, AGU Geophysical Monograph Series; 1–9.
- Roscoe JH. 1954. Ice streams. *Polar Record* **7**: 231–232.
- Rose KE. 1979. Characteristics of ice flow in Marie Byrd Land. *Journal of Glaciology* **24**(90): 63–75.
- Ross M, Campbell JE, Parent M, Adams RS. 2009. Palaeo-ice streams and the subglacial landscape mosaic of the North American mid-continental prairies. *Boreas* **38**(3): 421–439.
- Schoof C. 2002. Basal perturbations under ice streams: form drag and surface expression. *Journal of Glaciology* **48**(162): 407–416.
- Schoof C. 2007. Pressure-dependent viscosity and interfacial instability in coupled ice-sediment flow. *Journal of Fluid Mechanics* **570**: 227–252.
- Schoof C, Clarke GKC. 2008. A model of spiral flows in basal ice and the formation of subglacial flutes based on a Reiner–Rivlin rheology for glacial ice. *Journal of Geophysical Research* **113**: B05204.
- Schroeder DM, Blankenship DD, Young DA. 2013. Evidence for a water system transition beneath Thwaites Glacier, West Antarctica. *Proceedings of the National Academy of Science USA* **110**(30): 12 225–12 228.
- Schroeder DM, Blankenship DD, Young DA, Witus AE, Anderson JB. 2014. Airborne radar sounding evidence for deformable sediments and outcropping bedrock beneath Thwaites Glacier, West Antarctica. *Geophysical Research Letters* **41**. <https://doi.org/10.1002/2014GL061645>.
- Sergienko OV, Creyts TT, Hindmarsh RCA. 2014. Similarity of organized patterns in driving and basal stresses of Antarctic and Greenland ice sheets beneath extensive areas of basal sliding. *Geophysical Research Letters* **41**. <https://doi.org/10.1002/2014GL059976>.
- Sergienko OV, Hindmarsh RCA. 2013. Regular patterns in frictional resistance of ice-stream beds seen by surface data inversion. *Science* **342**: 1086–1089.
- Sergienko OV, Hulbe CL. 2011. 'sticky spots' and subglacial lakes under ice streams of the Siple Coast, Antarctica. *Annals of Glaciology* **52**(58): 18–22.
- Shabtaie S, Bentley CR. 1987. West Antarctic ice streams draining into the Ross Ice Shelf: configuration and mass balance. *Journal of Geophysical Research* **92**: 1311–1336.
- Shabtaie S, Whillans IM, Bentley CR. 1987. The morphology of ice streams A, B and C, West Antarctica, and their environs. *Journal of Geophysical Research* **92**(B9): 8865–8883.
- Shaw J. 1983. Drumlin formation related to inverted melt-water erosional marks. *Journal of Glaciology* **29**: 185–214.
- Shaw J, Faragini DM, Kvill DR, Rains BR. 2000. The Athabasca fluting field, Alberta, Canada: implications for the formation of large-scale fluting (erosional lineations). *Quaternary Science Reviews* **19**(10): 959–980.
- Shaw J, Piper DJW, Fader GBJ, King EL, Todd BJ, Bell T, Batterson MJ, Liverman DGE. 2006. A conceptual model of the deglaciation of Atlantic Canada. *Quaternary Science Reviews* **25**(17–18): 2059–2081.
- Shaw J, Pugin A, Young RR. 2008. A meltwater origin for Antarctic shelf bedforms with special attention to megalineations. *Geomorphology* **75**: 157–171.
- Shaw J, Young RR. 2010. Reply to comment by Ó Cofaigh, Dowdeswell, King, Anderson, Clark, DJA Evans, J. Evans, Hindmarsh, Lardner [sic] and Stokes 'Comments on Shaw J, Pugin A, Young R. (2009): a meltwater origin for Antarctic Shelf bedforms with special attention to megalineations'. *Geomorphology* **102**: 364–375. *Geomorphology* **117**: 199–201.
- Shepherd A, Ivins ER, Gerou A, Barletta VR, Bentley MJ, Bettadpur S, Briggs KH, Bromwich DH, Forsberg R, Galin N, Horwath M, Jacobs S, Joughin I, King MA, Lenaerts JTM, Li J, Ligtenberg SRM, Luckman A, Luthcke SB, McMillan M, Meister R, Milne G, Mougnot J, Muir A, Nicolas JP, Paden J, Payne AJ, Pritchard H, Rignot E, Rott H, Sorensen LS, Scambos TA, Scheuchl B, Schrama EJO, Smith B, Sundal AV, van Angelen JH, van de Berg WJ, van den Broeke MR, Vaughan DG, Velicogna I, Wahr J, Whitehouse PL, Wingham DJ, Yi D, Young D, Zwally HJ. 2012. A reconciled estimate of ice sheet mass balance. *Science* **338**: 1183–1189.
- Shipp S, Anderson J, Domack E. 1999. Late Pleistocene-Holocene retreat of the West Antarctic Ice Sheet system in the Ross Sea: Part

- 1 – Geophysical results. *Geological Survey of America, Bulletin* **111**(10): 1486–1516.
- Shipp SS, Wellner JS, Anderson JB. 2002. Retreat signature of a polar ice stream: sub-glacial geomorphic features and sediments from the Ross Sea, Antarctica. In *Glacier-influenced Sedimentation on High-Latitude Continental Margins*, Dowdeswell JA, Ó Cofaigh C (eds). London, Special Publication: Geological Society; 277–304.
- Siegert MJ, Carter S, Tabacco I, Popov S, Blankenship DD. 2005. A revised inventory of Antarctic subglacial lakes. *Antarctic Science* **17**: 453–460.
- Siegert MJ, Taylor J, Payne AJ, Hubbard B. 2004. Macro-scale bed roughness of the Siple Coast ice streams in West Antarctica. *Earth Surface Processes and Landforms* **29**: 1591–1596.
- Slater G. 1926. Glacial tectonics as reflected in disturbed drift deposits. *Proceedings of the Geologists Association* **37**(4): 392–400.
- Smith AM. 1997a. Basal conditions on Rutford Ice Stream, West Antarctica, from seismic observations. *Journal of Geophysical Research* **102**(B1): 543–552.
- Smith AM. 1997b. Variations in basal conditions on Rutford Ice Stream, West Antarctica. *Journal of Glaciology* **43**(144): 245–255.
- Smith AM. 2006. Microearthquakes and subglacial conditions. *Geophysical Research Letters* **33**: L24501. <https://doi.org/10.1029/2006GL028207>.
- Smith AM, Bentley CR, Bingham RG, Jordan TA. 2012. Rapid subglacial erosion beneath Pine Island Glacier, West Antarctica. *Geophysical Research Letters* **39**: L12501, <https://doi.org/10.1029/2012GL051651>.
- Smith AM, Jordan TA, Ferracciolo F, Bingham RG. 2013. Influence of subglacial conditions on ice stream dynamics: seismic and potential field data from Pine Island Glacier, West Antarctica. *Journal of Geophysical Research - Earth Surface* **118**: 1471–1482.
- Smith AM, Murray T. 2009. Bedform topography and basal conditions beneath a fast-flowing West Antarctic ice stream. *Quaternary Science Reviews* **28**: 584–596.
- Smith AM, Murray T, Nicholls KW, Makinson K, Algeirsdóttir G, Behar AE, Vaughan DG. 2007. Rapid erosion, drumlin formation, and changing hydrology beneath an Antarctic ice stream. *Geology* **35**(2): 127–130.
- Smith BE, Fricker HA, Joughin IR, Tulaczyk S. 2009. An inventory of active subglacial lakes in Antarctica detected by ICESat (2003–2008). *Journal of Glaciology* **55**(192): 573–595.
- Smith EC, Smith AM, White RS, Brisbourne AM, Pritchard HD. 2015. Mapping the ice–bed interface characteristics of Rutford Ice Stream, West Antarctica, using microseismicity. *Journal of Geophysical Research - Earth Surface* **120**: 1881–1894.
- Smith HTU. 1948. Giant glacial grooves. *American Journal of Science* **246**: 503–514.
- Smith JA, Andersen TJ, Shortt M, Gaffney AM, Truffer M, Stanton TP, Bindschadler R, Dutrieux P, Jenkins A, Hillenbrand C-D, Ehrmann W, Corr HFJ, Farley N, Crowhurst S, Vaughan DG. 2017. Sub-ice-shelf sediments record twentieth-century retreat of Pine Island glacier. *Nature* **541**: 77–80.
- Spagnolo M, Bartholomaeus TC, Clark CD, Stokes CR, Atkinson N, Dowdeswell JA, Ely JC, Graham A, Hogan KA, King E, Livingstone SJ, Pritchard HD. 2017. The periodic topography of ice stream beds: insights from the Fourier spectra of mega-scale glacial lineations. *Journal of Geophysical Research - Earth Surface* **122**: 1355–1373. <https://doi.org/10.1002/2016JF004154>
- Spagnolo M, Clark CD, Ely JC, Stokes CR, Andreassen K, Graham AGC, King EC. 2014. Size, shape and spatial arrangements of mega-scale glacial lineations from a large and diverse dataset. *Earth Surface Processes and Landforms* **39**: 1432–1448.
- Spagnolo M, Phillips E, Piotrowski JA, Rea BR, Clark CD, Stokes CR, Carr SJ, Ely JC, Ribolini A, Wysota W, Szuman I. 2016. Ice stream motion facilitated by a shallow-deforming and accreting bed. *Nature Communications* **7**: 10723. <https://doi.org/10.1038/ncomms10723>.
- Stanford SD, Mickelson DM. 1985. Till fabric and deformational structures in drumlins near Waukesha, Wisconsin, USA. *Journal of Glaciology* **31**(109): 220–228.
- Stearns LA, Smith BE, Hamilton GS. 2008. Increased flow speed on a large East Antarctic outlet glacier caused by subglacial floods. *Nature Geoscience* **1**: 827–831.
- Stephenson SN, Doake CSM. 1982. Dynamic behaviour of Rutford Ice Stream. *Annals of Glaciology* **3**: 295–299.
- Stokes CR. 2017. Deglaciation of the Laurentide Ice Sheet from the Last Glacial Maximum. *Cuadernos de Investigación Geográfica* **43**(2): 377–428.
- Stokes CR, Clark CD. 1999. Geomorphological criteria for identifying Pleistocene ice streams. *Annals of Glaciology* **28**: 67–74.
- Stokes CR, Clark CD. 2001. Palaeo-ice streams. *Quaternary Science Reviews* **20**: 1437–1457.
- Stokes CR, Clark CD. 2002a. Are long subglacial bedforms indicative of fast ice flow? *Boreas* **31**: 239–249.
- Stokes CR, Clark CD. 2002b. Ice stream shear margin moraines. *Earth Surface Processes and Landforms* **27**: 547–558.
- Stokes CR, Clark CD. 2003. The Dubawnt Lake palaeo-ice stream: evidence for dynamic ice sheet behaviour on the Canadian Shield and insights regarding the controls on ice stream location and vigour. *Boreas* **32**: 263–279.
- Stokes CR, Clark CD, Lian O, Tulaczyk S. 2006. Geomorphological map of ribbed moraine on the Dubawnt Lake Ice Stream bed: a signature of ice stream shut-down? *Journal of Maps* **2006**: 1–9.
- Stokes CR, Clark CD, Lian OB, Tulaczyk S. 2007. Ice stream sticky spots: a review of their identification and influence beneath contemporary and palaeo-ice streams. *Earth-Science Reviews* **81**: 217–249.
- Stokes CR, Fowler AC, Clark CD, Hindmarsh RCA, Spagnolo M. 2013a. The instability theory of drumlin formation and its explanation of their varied composition and internal structure. *Quaternary Science Reviews* **62**: 77–96.
- Stokes CR, Lian OB, Tulaczyk S, Clark CD. 2008. Superimposition of ribbed moraines on a palaeo-ice stream bed: implications for ice stream dynamics and shut-down. *Earth Surface Processes and Landforms* **33**(4): 593–609.
- Stokes CR, Margold M, Clark CD, Tarasov L. 2016a. Ice stream activity scaled to ice sheet volume during Laurentide ice sheet deglaciation. *Nature* **530**: 322–326.
- Stokes CR, Margold M, Creyts TT. 2016b. Ribbed bedforms on palaeo-ice stream beds resemble regular patterns of basal shear stress ('traction ribs') inferred from modern ice streams. *Journal of Glaciology* **62**(234): 696–713.
- Stokes CR, Spagnolo M, Clark CD, Ó Cofaigh C, Lian OB, Dunstone RB. 2013b. Formation of mega-scale glacial lineations on the Dubawnt Lake Ice Stream bed: 1. Size, shape and spacing from a large remote sensing dataset. *Quaternary Science Reviews* **77**: 190–209.
- Stokes CR, Tarasov L. 2010. Ice streaming in the Laurentide Ice Sheet: a first comparison between data-calibrated numerical model output and geological evidence. *Geophysical Research Letters* **37**: L01501. <https://doi.org/10.1029/2009GL040990>.
- Swithbank CWM. 1954. Ice streams. *Polar Record* **7**: 185–186.
- Taylor J, Siegert MJ, Payne AJ, Hubbard B. 2004. Regional-scale bed roughness beneath ice masses: measurement and analysis. *Computers and Geosciences* **30**: 899–908.
- Thomas RH. 1976. Thickening of the Ross Ice Shelf and equilibrium state of the West Antarctic Ice Sheet. *Nature* **259**(5540): 180–183.
- Thomas RH, Sanderson TJO, Rose KE. 1979. Effect of climate warming on the West Antarctic ice sheet. *Nature* **277**: 355–358.
- Thorsteinsson T, Raymond CF. 2000. Sliding versus till deformation in the fast motion of an ice stream over a viscous till. *Journal of Glaciology* **46**(155): 633–640.
- Trommelen MS, Ross M. 2014. Distribution and type of sticky spots at the centre of a deglacial streamlined lobe in northeastern Manitoba, Canada. *Boreas* **43**(3): 557–576.
- Trommelen MS, Ross M, Ismail A. 2014. Ribbed moraines in northern Manitoba, Canada: characteristics and preservation as part of a subglacial bed mosaic near the core regions of ice sheets. *Quaternary Science Reviews* **87**: 135–155.
- Truffer M, Echelmeyer KA. 2003. Of isbræ and ice streams. *Annals of Glaciology* **36**: 66–72.
- Tulaczyk SM. 2006. Scale independence of till rheology. *Journal of Glaciology* **52**(178): 377–380.
- Tulaczyk SM, Kamb B, Engelhardt HF. 2000a. Basal mechanics of Ice Stream B. I. Till mechanics. *Journal of Geophysical Research* **105**: 463–481.
- Tulaczyk SM, Kamb B, Engelhardt HF. 2000b. Basal mechanics of Ice Stream B. II. Plastic-undrained bed model. *Journal of Geophysical Research* **105**: 483–494.

- Tulaczyk SM, Kamb B, Scherer R, Engelhardt HF. 1998. Sedimentary processes at the base of a West Antarctica ice stream: constraints from textural and compositional properties of subglacial debris. *Journal of Sedimentary Research* **68**: 487–496.
- Tulaczyk SM, Scherer RP, Clark CD. 2001. A ploughing model for the origin of weak tills beneath ice streams: a qualitative treatment. *Quaternary International* **86**: 59–70.
- Van den Broeke M, Bamber J, Ettema J, Rignot E, Schrama E, van de Berg W, van Maijgaard E, Velicogna I, Wouters B. 2009. Partitioning recent Greenland mass loss. *Science* **326**: 984–986.
- Vornberger PL, Whillans IM. 1986. Surface features of Ice Stream B, Marie Byrd Land, West Antarctica. *Annals of Glaciology* **8**: 168–170.
- Vorren TO, Laberg JS. 1997. Trough mouth fans – palaeoclimate and ice-sheet monitors. *Quaternary Science Reviews* **16**: 865–881.
- Vorren TO, Laberg JS, Blaumme F, Dowdeswell JA, Kenyon NH, Mienert J, Rumohr J, Werner F. 1998. The Norwegian-Greenland Sea continental margins: Morphology and Late Quaternary sedimentary processes and environments. *Quaternary Science Reviews* **17**(1–3): 273–302.
- Walder JS, Fowler A. 1994. Channelized subglacial drainage over a deformable bed. *Journal of Glaciology* **40**(134): 3–15.
- Weertman J. 1969. Discussion: ‘What are glaciers surges?’ by Meier and Post (1969). *Canadian Journal of Earth Sciences* **4**(2): 816–817.
- Weertman J. 1974. Stability of the junction of an ice sheet and an ice shelf. *Journal of Glaciology* **13**(67): 3–11.
- Weertman J. 1976. Glaciology’s grand unsolved problem. *Nature* **260**(5549): 284–286.
- Wellner JS, Lowe AL, Shipp SS, Anderson JB. 2001. Distribution of glacial geomorphic features on the Antarctic continental shelf and correlation with substrate: implications for ice behaviour. *Journal of Glaciology* **47**(158): 397–411.
- Whillans IM. 1976. Radio-echo layers and the recent stability of the West Antarctic ice sheet. *Nature* **264**(5582): 125–155.
- Whillans IM, Bolzan J, Shabtaie S. 1987. Velocity of ice streams B and C, Antarctica. *Journal of Geophysical Research* **92**(B9): 8895–8902.
- Wingham DJ, Siegert MJ, Shepherd A. 2006. Muir AS. *Rapid discharge connects Antarctic lakes: Nature* **440**: 1033–1036.
- Winsborrow M, Andreassen K, Hubbard A, Plaze-Faverola A, Gudlaugsson E, Patton H. 2016. Regulation of ice stream flow through subglacial formation of gas hydrates. *Nature Geoscience* **9**: 370–374.
- Wright HE, Jr. 1973. Tunnel valleys, glacial surges, and subglacial hydrology of the Superior lobe, Minnesota. *Geological Society of America Memoirs* **316**: 251–276.